

PETERSON ISOMORPHISM IN K -THEORY AND RELATIVISTIC TODA LATTICE

ABSTRACT. The K -homology ring of the affine Grassmannian of $SL_n(\mathbb{C})$ was studied by Lam, Schilling, and Shimozono. It is realized as a certain concrete Hopf subring of the ring of symmetric functions. On the other hand, for the quantum K -theory of the flag variety Fl_n , Kirillov and Maeno provided a conjectural presentation based on the results obtained by Givental and Lee. We construct an explicit birational morphism between the spectrums of these two rings. Our method relies on Ruijsenaars's relativistic Toda lattice with unipotent initial condition. From this result, we obtain a K -theory analogue of the so-called Peterson isomorphism for (co)homology. We provide a conjecture on the detailed relationship between the Schubert bases, and, in particular, we determine the image of Lenart–Maeno's quantum Grothendieck polynomial associated with a Grassmannian permutation.

1. INTRODUCTION

Let Fl_n be the variety of complete flags $V_\bullet = (V_1 \subset \cdots \subset V_n = \mathbb{C}^n)$ in $\mathbb{C}^n$, which is a homogeneous space G/B , where $G = SL_n(\mathbb{C})$ and B is the Borel subgroup of the upper triangular matrices in G . Let $K(Fl_n)$ be the Grothendieck ring of coherent sheaves on Fl_n . Givental and Lee [16] studied the quantum K -theory $QK(Fl_n)$, which is a ring defined as a deformation of $K(Fl_n)$ (see [14, 37] for general construction of quantum K -theory). Similar to Givental–Kim's presentation (see [15, 20]) for the quantum cohomology ring $QH^*(Fl_n)$, Kirillov and Maeno [21] provided a conjectural presentation for $QK(Fl_n)$, which we denote temporarily by $\mathcal{QK}(Fl_n)$. Let $Gr_{SL_n} = G(\mathbb{C}((t)))/G(\mathbb{C}[[t]])$ be the *affine Grassmannian* of $G = SL_n$, whose K -homology $K_*(Gr_{SL_n})$ has a natural structure of a Hopf algebra. Lam, Schilling, and Shimozono [28] constructed a Hopf isomorphism between $K_*(Gr_{SL_n})$ and the subring $\Lambda_{(n)} := \mathbb{C}[h_1, \dots, h_{n-1}]$ of the ring Λ of symmetric functions. The first main result of this paper is an explicit ring isomorphism between $K_*(Gr_{SL_n})$ and $\mathcal{QK}(Fl_n)$ after appropriate localization.

The corresponding result in (co)homology, for a semisimple linear algebraic group G , is called the *Peterson isomorphism*; This result was presented in lectures by Peterson in MIT, 1997, and published in a paper by Lam and Shimozono [30] on torus-equivariant and parabolic settings.

We also provide a conjecture that describes a detailed correspondence between the Schubert bases for $K_*(Gr_{SL_n})$ and $\mathcal{QK}(Fl_n)$ (Conjecture 2).

Our method of constructing the isomorphism relies on Ruijsenaars's relativistic Toda lattice [45]. Note that a similar approach to the original Peterson isomorphism for SL_n by solving the non-relativistic Toda lattice was given by Lam and Shimozono [29, 31].

It is natural to ask how K -theoretic Peterson isomorphisms for a general semisimple linear algebraic group G should be constructed. There are some results indicating that an approach to this problem using integrable systems would be fruitful. The relativistic Toda lattice associated with any root system was introduced by

Kruglinskaya and Marshakov in [24]. The commuting family of q -difference Toda operators given by Etingof, as well as general such operators constructed by using a quantized enveloping algebra $U_q(\mathfrak{g})$ of a complex semisimple Lie algebra $\mathfrak{g}$, were discussed by Givental and Lee [16]. As for K -homology of the affine Grassmannian: Bezrukavnikov, Finkelberg, and Mirković [2] showed that the spectrum of the $G(\mathbb{C}[[t]])$ -equivariant K -homology ring of Gr_G is naturally identified with the *universal centralizer* of the Langlands dual group of G .

1.1. Relativistic Toda lattice. The *relativistic Toda lattice*, introduced by Ruijsenaars [45], is a completely integrable Hamiltonian system. This system can be viewed as an isospectral deformation of the Lax matrix $L = AB^{-1}$ with

$$(1.1) \quad A = \begin{pmatrix} z_1 & -1 & & & \\ & z_2 & -1 & & \\ & & \ddots & \ddots & \\ & & & z_{n-1} & -1 \\ & & & & z_n \end{pmatrix}, \quad B = \begin{pmatrix} 1 & & & & \\ -Q_1 z_1 & 1 & & & \\ & \ddots & \ddots & & \\ & & \ddots & 1 & \\ & & & -Q_{n-1} z_{n-1} & 1 \end{pmatrix},$$

where z_i and Q_j are complex numbers. The relativistic Toda lattice is a partial differential equation with independent variables $t_1, \dots, t_{n-1}$ expressed in the Lax form:

$$(1.2) \quad \frac{dL}{dt_i} = [L, (L^i)_<] \quad (i = 1, \dots, n-1),$$

where $(L^i)_<$ denotes the strictly lower triangular part of L^i . Consider the characteristic polynomial of L :

$$\Psi_L(\zeta) = \det(\zeta \cdot 1 - L) = \zeta^n + \sum_{i=1}^n (-1)^i F_i(z, Q) \zeta^{n-i}.$$

More explicitly, we have

$$(1.3) \quad F_i(z_1, \dots, z_n, Q_1, \dots, Q_{n-1}) = \sum_{\substack{I \subset \{1, \dots, n\} \\ \#I=i}} \prod_{j \in I} z_j \prod_{j \in I, j+1 \notin I} (1 - Q_j),$$

where $Q_n := 0$. Let $\gamma = (\gamma_1, \dots, \gamma_n) \in \mathbb{C}^n$. In this paper, we assume $\gamma_n = 1$ and consider the *isospectral variety*

$$(1.4) \quad Z_\gamma := \{(z, Q) \in \mathbb{C}^{2n-1} \mid F_i(z, Q) = \gamma_i \ (1 \leq i \leq n)\}.$$

This is an affine algebraic variety with coordinate ring

$$\mathbb{C}[Z_\gamma] = \mathbb{C}[z_1, \dots, z_n, Q_1, \dots, Q_{n-1}] / I_\gamma, \quad I_\gamma = \langle F_i(z, Q) - \gamma_i \ (1 \leq i \leq n) \rangle.$$

Note that Z_γ is isomorphic to a locally closed subvariety of $SL_n(\mathbb{C})$ (see §3.2).

1.2. Quantum K -theory of flag variety. A particularly interesting case occurs when

$$\gamma_i = \binom{n}{i} \quad (1 \leq i \leq n).$$

Then, the corresponding Lax matrix L has characteristic polynomial $(\zeta - 1)^n$. In fact, in this case, L is principal unipotent in the sense that it has only one Jordan block such that the eigenvalues are all 1, so we say the corresponding isospectral variety is *unipotent* and denote it by Z_{uni} . Let $\mathcal{L}_i$ be the *tautological line bundle* whose fiber over $V_\bullet \in Fl_n$ is V_i/V_{i-1} .

Conjecture 1.1 (Kirillov–Maeno [21]). *There is a canonical ring isomorphism*

$$(1.5) \quad QK(Fl_n) \simeq \mathbb{C}[Z_{\text{uni}}],$$

with z_i identified with the class the tautological line bundle $\mathcal{L}_i$.

Givental and Lee [16] studied a certain generating function of the $\mathbb{C}^* \times SL_n(\mathbb{C})$ -equivariant Euler characteristic of a natural family of line bundles on the *quasimap spaces* from $\mathbb{P}^1$ to Fl_n . The function is a formal power series in $Q = (Q_1, \dots, Q_{n-1})$ and its copy Q' , depending on q , the coordinate of $\mathbb{C}^*$, and $(\Lambda_1, \dots, \Lambda_n)$ with $\Lambda_1 \cdots \Lambda_n = 1$, the coordinates of the maximal torus of $SL_n(\mathbb{C})$, respectively. They proved that the generating function is the eigenfunction of the finite q -difference Toda operator. A relation to the Toda lattice has been studied further by Braverman and Finkelberg [3, 4]. Accordingly, it has been expected that there is a finite q -difference counterpart of quantum D -module ([12, 13], see also [20]) giving the structure of $QK(Fl_n)$; however, to the best of the authors' knowledge, the connection between the multiplication in $QK(Fl_n)$ and the q -difference system is still uncertain. We hope that recent work by Iritani, Milanov, and Tonita [Iritani et al.(2015)] gives an explanation of this connection, and that ultimately Conjecture 1.1 will be proved.

We add some remarks on a certain finiteness property of the quantum K -theory. Note that the ring is originally defined as a $\mathbb{C}[[Q]]$ -algebra (see [14, 37]). For cominuscule G/P , such as the Grassmannian $Gr_d(\mathbb{C}^n)$ of d -dimensional subspaces of $\mathbb{C}^n$, it was shown by Buch, Chaput, Mihalcea, and Perrin [7, 8] that multiplicative structure constants for the (quantum) Schubert basis $[\mathcal{O}_{X_w}]$ are polynomial in Q_i (see also Buch and Mihalcea [9] for earlier results on the Grassmannian). Thus, for such a variety, the $\mathbb{C}[Q]$ -span of the Schubert classes forms a subring. So far, it is not known whether or not the finiteness holds for Fl_n in general. In the above conjecture, $QK(Fl_n)$ should be interpreted as the $\mathbb{C}[Q]$ -span of the Schubert classes.

With this conjecture in mind, in this paper, we will denote $\mathbb{C}[Z_{\text{uni}}]$ by $\mathcal{QK}(Fl_n)$ in distinction to $QK(Fl_n)$.

1.3. Dual stable Grothendieck polynomials. Let Λ denote the complexified¹ ring of symmetric functions (see [40]). If we denote by h_i the i th *complete symmetric function*, Λ is the polynomial ring $\mathbb{C}[h_1, h_2, \dots]$. It has the so-called *Hall inner product* $\langle \cdot, \cdot \rangle : \Lambda \times \Lambda \rightarrow \mathbb{C}$ and a standard Hopf algebra structure.

For each partition $\lambda = (\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_\ell)$ of length ℓ , the *Schur functions* $s_\lambda \in \Lambda$ are defined by

$$(1.6) \quad s_\lambda = \det(h_{\lambda_i + j - i})_{1 \leq i, j \leq \ell}.$$

The *stable Grothendieck polynomial* G_λ is given as the sum of *set-valued tableaux* of shape λ , and they form a basis of a completed ring $\hat{\Lambda}$ of symmetric functions (see Buch [5] for details). The *dual stable Grothendieck polynomials* $\{g_\lambda\}$ due to Lam and Pylyavskyy [27] are defined by $\langle g_\lambda, G_\mu \rangle = \delta_{\lambda, \mu}$. It was shown in [27] that $\{g_\lambda\}$ is identified with the K -homology Schubert basis of the infinite Grassmannian (see [27, §9.5] for a more precise statement). Shimozono and Zabrocki [46] proved

¹It is possible to work over the integers; however, we use complex coefficients throughout the paper because we use many coordinate rings of complex algebraic varieties.

a determinant formula for g_λ . The following formula for g_λ is also available:

$$(1.7) \quad g_\lambda = \det \left(\sum_{m=0}^{\infty} (-1)^m \binom{1-i}{m} h_{\lambda_i+j-i-m} \right)_{1 \leq i, j \leq \ell} = s_\lambda + \text{lower degree terms},$$

where $h_m = 0$ for $m < 0$.

Remark 1.2. *It is straightforward to see the equivalence of formula (1.7) and the one in [46] (see also [35, Equation (4)]), so we omit the details.*

1.4. Lam–Schilling–Shimozono’s presentation for $K_*(Gr_{SL_n})$. In [28], Lam, Schilling and Shimozono showed that the K -homology of the affine Grassmannian $K_*(Gr_{SL_n})$ [28, §5.1] can be realized as a subring in the affine K -theoretic nil-Hecke algebra of Kostant–Kumar [23]. Let us denote by $\Lambda_{(n)} := \mathbb{C}[h_1, \dots, h_{n-1}]$ the subring of Λ generated by $h_1, \dots, h_{n-1}$. In [28], the following ring isomorphism was established:

$$(1.8) \quad K_*(Gr_{SL_n}) \simeq \Lambda_{(n)}.$$

Note that $K_*(Gr_{SL_n})$ is equipped with a Hopf algebra structure coming from the based loop space $\Omega SU(n)$, and the above is a Hopf isomorphism with the Hopf algebra structure on $\Lambda_{(n)}$ induced from the canonical one on Λ .

1.5. K -theoretic Peterson isomorphism. Let Z_{uni}° be the Zariski open set of Z_{uni} defined as the complement of the divisor defined by $Q_1 \cdots Q_{n-1} = 0$. Thus the coordinate ring $\mathbb{C}[Z_{\text{uni}}^\circ]$ is a localization of $\mathbb{C}[Z_{\text{uni}}] = \mathbb{C}[z, Q]/I_{\text{uni}}$ by Q_i ($1 \leq i \leq n-1$). In view of Conjecture 1.1, we define

$$(1.9) \quad \mathcal{QK}(Fl_n)_{\text{loc}} := \mathbb{C}[Z_{\text{uni}}^\circ].$$

For the affine Grassmannian side, we define

$$(1.10) \quad K_*(Gr_{SL_n})_{\text{loc}} := \Lambda_{(n)}[\sigma_i^{-1}, \tau_i^{-1} \ (1 \leq i \leq n-1)],$$

where

$$(1.11) \quad \tau_i = g_{R_i}, \quad \sigma_i = \sum_{\mu \subset R_i} g_\mu \quad (1 \leq i \leq n-1),$$

where R_i is the rectangle² with i rows and $(n-i)$ columns. We set $\tau_0 = \sigma_0 = \tau_n = \sigma_n = 1$.

Example 1.3. *Let $n = 3$. Then we have*

$$\tau_1 = h_2, \quad \tau_2 = h_1^2 - h_2 + h_1, \quad \sigma_1 = h_2 + h_1 + 1, \quad \sigma_2 = h_1^2 - h_2 + 2h_1 + 1.$$

The main result of this paper is the following theorem.

Theorem 1.4 (Corollary 2.5, Propositions 3.10, 4.2, 4.5). *There is an isomorphism of rings*

$$(1.12) \quad \Phi_n : \mathcal{QK}(Fl_n)_{\text{loc}} \xrightarrow{\sim} K_*(Gr_{SL_n})_{\text{loc}},$$

given by

$$(1.13) \quad z_i \mapsto \frac{\tau_i \sigma_{i-1}}{\sigma_i \tau_{i-1}} \quad (1 \leq i \leq n), \quad Q_i \mapsto \frac{\tau_{i-1} \tau_{i+1}}{\tau_i^2} \quad (1 \leq i \leq n-1).$$

²Our notation of R_i is conjugate (transpose) to the one used in [31].

Remark 1.5. Let p_i be the i th power sum symmetric function in Λ . With the identification $t_i = p_i/i$ ($1 \leq i \leq n-1$), the formulas (1.11), (1.12) give an explicit solution of the relativistic Toda lattice that is a rational function in time variables $t_1, \dots, t_{n-1}$. Thus, if we substitute the rational expression of (z, Q) in t_j into $F_i(z, Q)$, then such a complicated rational function is equal to $\binom{n}{i}$. This remarkable identity can be understood as a consequence of the isospectral property of the relativistic Toda lattice.

1.6. Method of construction. The ring homomorphism Φ_n is obtained by solving the relativistic Toda lattice. Our method of solving the system is analogous to the one employed by Kostant [22] for the finite non-periodic Toda lattice. Let C_γ be the companion matrix of the (common) characteristic polynomial of $L \in Z_\gamma$. We denote the centralizer of C_γ by $\mathfrak{X}_\gamma$, which is the affine space of dimension n . We can construct a birational morphism α from Z_γ to $\mathbb{P}(\mathfrak{X}_\gamma)$. On the open set of $\mathbb{P}(\mathfrak{X}_\gamma)$ defined as the complement of the divisors given by τ_i, σ_i , we can construct the inverse morphism β to α . When γ is unipotent, the associated isomorphism between the coordinate rings of these open sets is nothing but Φ_n .

1.7. Quantum Grothendieck polynomials. The *quantum Grothendieck polynomials* $\mathfrak{G}_w^Q$ [38] of Lenart and Maeno are a family of polynomials in the variables $x_1, \dots, x_n$ and quantum parameters $Q_1, \dots, Q_{n-1}$ indexed by permutations $w \in S_n$ (see §5.2 for the definition). Note that x_i is identified with $1 - z_i$ (the first Chern class of the dual line bundle of $\mathcal{L}_i$). It was conjectured in [38] that the polynomial $\mathfrak{G}_w^Q$ represents the quantum Schubert class $[\mathcal{O}_{X_w}]$ in $\mathbb{C}[z, Q]/I_{\text{uni}}$ under the conjectured isomorphism³ (1.5).

1.8. K -theoretic k -Schur functions. In the sequel we use the notation $k = n-1$. Let $\mathcal{B}_k$ denote the set of k -bounded partitions λ , i.e. the partitions such that $\lambda_1 \leq k$. The K -theoretic k -Schur functions $\{g_\lambda^{(k)}\} \subset \Lambda_{(n)} \simeq K_*(Gr_{SL_n})$ are indexed by $\lambda \in \mathcal{B}_k$ and defined in [28] as the dual basis of the Schubert basis of the Grothendieck ring $K^*(Gr_{SL_n})$ of the thick version of the affine Grassmannian (see [19] for Kashiwara's construction of the thick flag variety of a Kac-Moody Lie algebra). The highest degree component of $g_\lambda^{(k)}$ is the k -Schur function $s_\lambda^{(k)}$ introduced by Lapointe, Lascoux, and Morse [32] (see also [25] and references therein).

We note that $g_\lambda^{(k)}$ is equal to g_λ if k is sufficiently large [28, Remark 7.3]. Morse [43]⁴ conjectured that for a partition λ such that $\lambda \subset R_i$ for some $1 \leq i \leq k$, we have

$$(1.14) \quad g_\lambda^{(k)} = g_\lambda,$$

thus in particular, $g_{R_i}^{(k)} = g_{R_i} = \tau_i$. This conjecture is relevant for our considerations. Note that the counterpart of this conjecture for k -Schur functions was proved in [34]; that is, $s_\lambda^{(k)} = s_\lambda$ if $\lambda \subset R_i$ for some $1 \leq i \leq k$. In particular, we have $s_{R_i}^{(k)} = s_{R_i}$. It is worth remarking s_{R_i} arise as the “ τ -functions” of the finite non-periodic Toda lattice with nilpotent initial condition.

³The main result of [38] is a Monk-type formula for $\mathfrak{G}_w^Q$. It is worthwhile to emphasize that the formula is proved logically independent from Conjecture 1.1. In fact, the formula holds in the polynomial ring of x_i and Q_i .

⁴The conjecture after Property 47 in [43].

1.9. Image of the quantum Grothendieck polynomials. We are interested in the image of $\mathfrak{S}_w^Q$ by Φ_n .

Let $\text{Des}(w) = \{i \mid 1 \leq i \leq n-1, w(i) > w(i+1)\}$ denote the set of descents of $w \in S_n$. A permutation $w \in S_n$ is d -Grassmannian if $\text{Des}(w) = \{d\}$; such elements are in bijection with partition $\lambda = (\lambda_1, \dots, \lambda_d)$ such that $\lambda \subset R_d$. If $\lambda \subset R_d$, then the corresponding d -Grassmannian permutation $w_{\lambda,d}$ (see Remark 6.7) has length $|\lambda| = \sum_{i=1}^d \lambda_i$. Let $\lambda^\vee := (n-d-\lambda_d, \dots, n-d-\lambda_1)$.

Theorem 1.6 (cf. Corollary 5.7). *Let λ be a permutation such that $\lambda \subset R_d$. Then we have*

$$(1.15) \quad \Phi_n(\mathfrak{S}_{w_{\lambda,d}}^Q) = \frac{g_{\lambda^\vee}}{\tau_d}.$$

For a general permutation w , we will provide a conjecture on the image of $\mathfrak{S}_w^Q$. Let $\lambda : S_n \rightarrow \mathcal{B}_k$ be a map defined by Lam and Shimozono [31, §6] (see §7.1 below). In order to state the conjecture we also need an involution ω_k on $\mathcal{B}_k$, $\mu \mapsto \mu^{\omega_k}$, called the k -conjugate ([33]). The image of the map λ consists of elements in $\mathcal{B}_k$ that is k -irreducible, that is, those k -bounded partition $\mu = (1^{m_1} 2^{m_2} \dots (n-1)^{m_{n-1}})$ such that $m_i \leq k-i$ ($1 \leq i \leq n-1$). Let $\mathcal{B}_k^*$ denote the set of all k -irreducible k -bounded partitions. Note that $\mathcal{B}_k^*$ is preserved by k -conjugate. Let us denote by S_n^* the subset $\{w \in S_n \mid w(1) = 1\}$ of S_n . We know that λ gives a bijection from S_n^* to $\mathcal{B}_k^*$ (see §7.1 below).

Conjecture 1.7. *Let w be in S_n . There is a polynomial $\tilde{g}_w \in \Lambda_{(n)}$ such that*

$$(1.16) \quad \Phi_n(\mathfrak{S}_w^Q) = \frac{\tilde{g}_w}{\prod_{i \in \text{Des}(w)} \tau_i}.$$

Furthermore, $\tilde{g}_w$ satisfies the following properties:

(i) If $\lambda(w) = \lambda(w')$ for $w, w' \in S_n$, then we have

$$\tilde{g}_w = \tilde{g}_{w'}.$$

(ii) For $w \in S_n$, we have

$$(1.17) \quad \tilde{g}_w = g_{\lambda(w)^{\omega_k}}^{(k)} + \sum_{\mu} a_{w,\mu} g_{\mu}^{(k)}, \quad a_{w,\mu} \in \mathbb{Z},$$

where μ runs for all elements in $\mathcal{B}_k^*$ such that $|\mu| < |\lambda(w)|$.

(iii) $(-1)^{|\mu| - |\lambda(w)|} a_{w,\mu}$ is a non-negative integer.

The counterpart of Conjecture 1.7 in the (co)homology case was established in [31], where the quantum Schubert polynomial $\mathfrak{S}_w^q$ of Fomin, Gelfand, and Postnikov [10] is sent by the original Peterson isomorphism to the fraction, whose numerator is the single k -Schur function associated with $\lambda(w)^{\omega_k}$, and the denominator is the products of $s_{R_i}^{(k)}$ such that $i \in \text{Des}(w)$. Note in the formula in [31], the numerator is the k -Schur function associated with $\lambda(w)$ without k -conjugate by reason of the convention.

If $w \in S_n$ is d -Grassmannian for some d , and $w = w_{\mu,d}$ with $\mu \subset R_d$, then from Theorem 1.6 we have

$$\tilde{g}_w = g_{\mu^\vee}.$$

Since we know that $\mu^\vee = \lambda(w_{\mu,d})^{\omega_k}$ (Lemma 7.1 below), if (1.14) is true, Conjecture 1.7 holds for all Grassmannian permutations w . In the early stage of this work, we

expected that $\tilde{g}_w$ is always a single K - k -Schur function, however, this is not the case; for example we have

$$\tilde{g}_{1423} = g_{2,1,1}^{(3)} - g_{2,1}^{(3)}.$$

1.10. Further discussions. Let us assume that Conjecture 1.7 is true and discuss its possible implications. The property (iii) says $\{\tilde{g}_w\}_{w \in S_n^*}$ and $\{g_\nu^{(k)}\}_{\mu \in \mathcal{B}_k^*}$ are two bases of the same space, and the transition matrix between the bases is lower unitriangular. This suggests that $\{\tilde{g}_w\}_{w \in S_n^*}$ is a part of an important basis of $\Lambda_{(n)}$ different from the K - k -Schur basis. One possibility of such basis will be the following: Let us denote the function $\tilde{g}_w$ by $\tilde{g}_\nu^{(k)}$ with $\nu = \lambda(w)^{\omega_k} \in \mathcal{B}_k^*$. For a general k -bounded partition μ , we can uniquely write it as $\mu = \nu \cup \bigcup_{i=1}^{n-1} R_i^{e_i}$, $\nu \in \mathcal{B}_k^*$, $e_i \geq 0$. Then we define

$$\tilde{g}_\mu^{(k)} = \tilde{g}_\nu^{(k)} \cdot \tau_1^{e_1} \cdots \tau_{n-1}^{e_{n-1}}.$$

One sees that $\tilde{g}_\mu^{(k)}$ ($\mu \in \mathcal{B}_k$) form a basis of $\Lambda_{(n)}$.

In the proof of the isomorphism in [30], they work in T -equivariant (T is the maximal torus of G) settings, and first give the module isomorphism ψ from $H_*^T(Gr_G)_{\text{loc}}$ to $QH_T^*(G/B)_{\text{loc}}$, and next prove the ψ -preimage of the quantum Chevalley formula. Since the quantum Chevalley formula uniquely characterizes $QH_T^*(G/B)$ due to a result of Mihalcea [41], we know that ψ is a ring isomorphism. In our situation, we proved the ring isomorphism Φ_n (Theorem 2.5) without using the quantum Monk formula of $\mathcal{QK}(Fl_n)$ (cf. Lenart-Postnikov [39]). Thus the basis $\{\tilde{g}_\nu^{(k)}\}_{\nu \in \mathcal{B}_k}$ should satisfy the corresponding formula in $\Lambda_{(n)}$. These issues will be studied further elsewhere.

1.11. Organization. In Sections 2–4 of this paper, we give the K -theoretic Peterson morphism and prove it is an isomorphism (Theorem 2.5). In Sections 5–6, we calculate the image of quantum Grothendieck polynomials associated with Grassmannian permutations (Theorem 1.6). In Section 7, we discuss some details of Conjecture 1.7.

In Section 2, we state the main results of the first main part of the paper. We construct a birational morphism α from Z_γ to $\mathbb{P}(\mathcal{O}_\gamma)$ with $\mathcal{O}_\gamma := \mathbb{C}[\zeta]/(\zeta^n + \sum_{i=1}^{n-1} (-1)^i \gamma_i \zeta^{n-i})$. We also describe the open sets of both Z_γ and $\mathbb{P}(\mathcal{O}_\gamma)$ that are isomorphic as affine algebraic varieties. The complement of the open part of $\mathbb{P}(\mathcal{O}_\gamma)$ is a divisor given by explicitly defined functions T_i, S_i . The main construction of Section 2 is the definition of the map Φ_n in Theorem 2.5. In fact, the isomorphism statement of Theorem 2.5 is given as Corollary 2.5, which is the unipotent case of Theorem 2.4. In Section 3, we prove Theorem 2.4. We also give a more conceptual description of the map α and its inverse β . Section 3.6 includes the formula of Φ_n in terms of the functions T_i, S_i . In Section 4 we determine the precise form of the τ -functions T_i, S_i in terms of dual stable Grothendieck polynomials, thus completing the proof of Theorem 2.5.

In Section 5, we summarize the second main result. The aim is to calculate the quantum Grothendieck polynomials associated with Grassmannian permutations. We give a version $\hat{Q}_d$ of the quantization map from the K -ring of the Grassmannian $Gr_d(\mathbb{C}^n)$ to $\mathcal{QK}(Fl_n)$ by using our K -Peterson isomorphism. The main statement of Section 5 is the compatibility of $\hat{Q}_d$ and Lenart–Maeno’s quantization map $\hat{Q}$ with respect to the embedding $K(Gr_d(\mathbb{C}^n)) \hookrightarrow K(Fl_n)$ (Theorem 5.6). As a corollary

to this, we obtain Theorem 1.6. Section 6 is devoted to the proof of Theorem 5.6. In Section 7, we explain some details of Conjecture 2 and give some examples of calculations.

2. CONSTRUCTION OF K -THEORETIC PETERSON ISOMORPHISM

In this section, we state our main construction.

Let γ_i ($1 \leq i \leq n-1$) be any complex numbers and set $\gamma_n = 1$. Let

$$f_\gamma(\zeta) = \zeta^n + \sum_{i=1}^n (-1)^i \gamma_i \zeta^{n-i}.$$

Let $\mathcal{O}_\gamma$ denote the quotient ring $\mathbb{C}[\zeta]/(f_\gamma(\zeta))$. We also consider $\mathcal{O}_\gamma$ as an affine space. For $L = AB^{-1} \in Z_\gamma$, let $\Delta_{i,j} = \Delta_{i,j}(z, Q)$ denote the (i, j) -minor determinant of $\zeta B - A$.

Definition 2.1. We define the map $\alpha : Z_\gamma \rightarrow \mathbb{P}(\mathcal{O}_\gamma)$ sending $L \in Z_\gamma$ to $[\Delta_{1,1}]$.

Example 2.2. If $n = 3$ then $\Delta_{1,1}$ is given as

$$\Delta_{1,1} = \zeta^2 + (Q_2 z_2 - z_2 - z_3)\zeta + z_2 z_3.$$

This is a handy definition of α . A more conceptual description of the map α is given in the next section, where $\mathcal{O}_\gamma$ is interpreted as the centralizer of the companion matrix C_γ of the characteristic polynomial f_γ . In fact, we will construct the inverse β of α defined on an open set of $\mathbb{P}(\mathcal{O}_\gamma)$, which is the counterpart of the map Kostant [22] defined for the ordinary finite Toda lattice.

The case of our interest, as was noted above, is $\gamma_i = \binom{n}{i}$ that is equivalent to $\Psi_L(\zeta) = (\zeta - 1)^n$. We call this parameter *unipotent* and denote the corresponding isospectral variety by Z_{uni} . Recall that we denote

$$(2.1) \quad \mathcal{QK}(Fl_n) = \mathbb{C}[Z_{\text{uni}}],$$

which is our working definition of the quantum K -theory of Fl_n .

We will show below that α is a birational morphism of algebraic varieties. We also describe open parts that are isomorphic via the map α explicitly. As the corresponding isomorphism between the coordinate rings, we obtain the K -theory analogue of the Peterson isomorphism.

Definition 2.3. Fix a linear isomorphism $\mathbf{c} : \mathcal{O}_\gamma \rightarrow \mathbb{C}^n$. For $0 \leq j \leq n$, and $\varphi \in \mathcal{O}_\gamma$, let $\mathbf{a}_j = \mathbf{c}(\zeta^j)$, $\mathbf{b}_j = \mathbf{c}(\varphi \zeta^j)$. Define for $1 \leq i \leq n$,

$$\begin{aligned} T_i(\varphi) &= |\mathbf{b}_0, \mathbf{b}_1, \dots, \mathbf{b}_{i-1}, \mathbf{a}_{i-1}, \dots, \mathbf{a}_{n-2}|, \\ S_i(\varphi) &= |\mathbf{b}_0, \mathbf{b}_1, \dots, \mathbf{b}_{i-1}, \mathbf{a}_i, \dots, \mathbf{a}_{n-1}|. \end{aligned}$$

Note that a different choice of $\mathbf{c}$ yields a change $T_i \mapsto c^i T_i$, $S_i \mapsto c^i S_i$ with a nonzero constant $c \in \mathbb{C}^*$. Such change does not effect the following constructions, however, we choose $\mathbf{c}$ so that $|\mathbf{a}_0, \dots, \mathbf{a}_{n-1}| = 1$. For each i , both T_i and S_i are homogenous polynomial functions in $\mathbb{C}[\mathcal{O}_\gamma]$ of degree i . Let $Y_\gamma = \mathbb{P}(\mathcal{O}_\gamma)$ and define a Zariski open set

$$(2.2) \quad Y_\gamma^\circ = \{[\varphi] \in \mathbb{P}(\mathcal{O}_\gamma) \mid T_i(\varphi) \neq 0, S_i(\varphi) \neq 0 \ (1 \leq i \leq n)\}.$$

Our first main result is the following.

Theorem 2.4. The map α gives an isomorphism from Z_γ° to Y_γ° as affine algebraic varieties.

Now we apply this to unipotent case, namely the case when $\gamma_i = \binom{n}{i}$. We choose $\mathbf{c} : \mathcal{O}_\gamma \rightarrow \mathbb{C}^n$, $\varphi \mapsto {}^t(c_0, c_1, \dots, c_{n-1})$ as

$$(2.3) \quad \varphi = \sum_{i=0}^{n-1} (-1)^i c_i \cdot (\zeta - 1)^i.$$

Then we have $T_n = S_n = c_0^n$. So Y_{uni}° is an open subvariety of the affine open set U_0 of $\mathbb{P}(\mathcal{O}_{\text{uni}})$ defined by $c_0 \neq 0$. We identify the coordinate ring $\mathbb{C}[c_1/c_0, \dots, c_{n-1}/c_0]$ of U_0 with $\Lambda_{(n)} = \mathbb{C}[h_1, \dots, h_{n-1}]$ by

$$(2.4) \quad h_i = c_i/c_0 \quad (1 \leq i \leq n-1).$$

Using this identification, we will prove (see §4)

$$(2.5) \quad \tau_i = T_i/c_0^i, \quad \sigma_i = S_i/c_0^i \quad (1 \leq i \leq n-1).$$

Via the isomorphism $K(\text{Gr}_{SL_n}) \simeq \Lambda_{(n)}$, we have $\mathbb{C}[Y_{\text{uni}}^\circ] = K(\text{Gr}_{SL_n})[\tau_i^{-1}, \sigma_i^{-1}]$.

Corollary 2.5. *We have the following isomorphism of rings:*

$$\Phi_n : \mathcal{QK}(Fl_n)[Q_i^{-1} (1 \leq i \leq n-1)] \xrightarrow{\sim} K(\text{Gr}_{SL_n})[\tau_i^{-1}, \sigma_i^{-1} (1 \leq i \leq n-1)].$$

The explicit formula of Φ_n (the second statement of Theorem 2.5) will be derived below in Proposition 3.10 in §3.6 together with Propositions 4.2 and 4.5.

3. PROOF OF THEOREM 2.4

This section is devoted to the proof of Theorem 2.4.

3.1. Gauss decomposition. We use the following notation of minor determinants for an $n \times n$ matrix $X = (x_{ij})_{1 \leq i, j \leq n}$:

$$(3.1) \quad \xi_{i_1, \dots, i_r}^{j_1, \dots, j_r}(X) = \det(x_{i_a, j_b})_{1 \leq a, b \leq r}.$$

Let $\mathbf{B}$ (resp. $\mathbf{B}_-$) denote the Borel subgroup of $GL_n(\mathbb{C})$ consisting of upper (resp. lower) triangular matrices. Let $\mathbf{N}_-$ (resp. $\mathbf{N}$) denote the subgroup consisting of the unipotent lower (resp. upper) triangular matrices.

Proposition 3.1. *A square matrix X of size n can be expressed as $X = X_+ \cdot X_-$ with $X_+ \in \mathbf{B}$, and $X_- \in \mathbf{N}_-$, if and only if $\xi_{i+1, \dots, n}^{i+1, \dots, n}(X) \neq 0$ for $1 \leq i \leq n-1$.*

Proof. This is the factorization known as the Gauss or the LU-decomposition. The result is standard. See [44] for example. $\square$

Let σ denote the matrix $\sum_{i=1}^{n-1} E_{i+1, i} + E_{1, n}$ which represents the cyclic permutation $(1, \dots, n)$. Let $\varepsilon := \text{diag}(1, -1, 1, \dots, (-1)^{n-1})$.

Proposition 3.2. *Let X be a square matrix n such that $x_{1, n} \neq 0$. Then X can be expressed as $X = U^{-1}R$ with $R = (r_{ij}) \in \mathbf{B}\sigma$, $U = (u_{ij}) \in \mathbf{N}_-\varepsilon$, if and only if $\xi_{1, \dots, i-1, i}^{1, \dots, i-1, n}(X) \neq 0$ for $2 \leq i \leq n-1$. Moreover, if such decomposition exists, we have*

$$(3.2) \quad r_{i+1, i} = (-1)^{i+1} \frac{\xi_{1, \dots, i, i+1}^{1, \dots, i, n}(X)}{\xi_{1, \dots, i-1, i}^{1, \dots, i-1, n}(X)} \quad (1 \leq i \leq n-1).$$

3.2. The variety Z of Lax matrices. Let $J = \sum_{i=1}^{n-1} E_{i,i+1}$. Let Z denote the set of matrices L in $SL_n(\mathbb{C})$ satisfying the following conditions:

- (Z₁) : $L + J$ is a lower triangular matrix,
- (Z₂) : all entries of L^{-1} further down the second subdiagonal are zero,
- (Z₃) : $\xi_{i+1,\dots,n}^{i+1,\dots,n}(L) \neq 0$ for $1 \leq i \leq n-1$.

Let T be the subgroup of $(\mathbb{C}^*)^n$ consisting of $(z_1, \dots, z_n)$ such that $z_1 \cdots z_n = 1$.

Proposition 3.3. *The map $T \times \mathbb{C}^{n-1} \rightarrow Z$ defined by sending (z, Q) with $z \in T$, $Q = (Q_1, \dots, Q_{n-1}) \in \mathbb{C}^{n-1}$ to $L = AB^{-1}$ with*

$$(3.3) \quad A = \sum_{i=1}^n z_i E_{i,i} - J, \quad B = 1 - \sum_{i=1}^{n-1} Q_i z_i E_{i+1,i},$$

is an isomorphism of algebraic varieties.

Proof. It is easy to see $L = AB^{-1}$ given by (3.3) satisfies (Z₁). L^{-1} is given as follows:

$$(3.4) \quad \begin{pmatrix} \frac{1}{z_1} & \frac{1}{z_1 z_2} & \cdots & \cdots & \frac{1}{z_1 z_2 \cdots z_n} \\ -Q_1 & -\frac{Q_1 - 1}{z_2} & \cdots & \cdots & -\frac{Q_1 - 1}{z_2 \cdots z_n} \\ 0 & -Q_2 & \ddots & \cdots & \vdots \\ 0 & 0 & \ddots & -\frac{Q_{n-2} - 1}{z_{n-1}} & -\frac{Q_{n-2} - 1}{z_{n-1} z_n} \\ 0 & 0 & 0 & -Q_{n-1} & -\frac{Q_{n-1} - 1}{z_n} \end{pmatrix},$$

and thereby (Z₂) holds. To see L satisfies (Z₃) we only need to notice that L is factorized as in Proposition 3.1. We construct the inverse map by using Proposition 3.1. Let $L \in Z$. We decompose it as $L = AB^{-1}$, where $A \in \mathbf{B}$, $B^{-1} \in \mathbf{N}_-$. Let $M = L^{-1}$. Define $Q_i = -M_{i+1,i}$ and $z_i = M_{1,i}/M_{1,i-1}$ with $M_{1,0} = 1$. It is straightforward, by using (Z₁) and (Z₂), to check A, B are given by (3.3). $\square$

Thus Z is the affine variety whose coordinate ring $\mathbb{C}[Z]$ is $\mathbb{C}[z, Q]/(z_1 \cdots z_n - 1)$. We define the subset Z° of Z by imposing the condition:

- (Z₄) : $Q_i \neq 0$ ($1 \leq i \leq n-1$).

Note that Z_γ defined by (1.4) is a closed subvariety of Z . Let $Z_\gamma^\circ = Z_\gamma \cap Z^\circ$.

3.3. Centralizer of C_γ . Let C_γ denote the companion matrix of $f_\gamma(\zeta)$. Explicitly $C_\gamma = J + \sum_{i=1}^n (-1)^{i-1} \gamma_i E_{n,n-i+1}$. Let $\mathfrak{X}_\gamma$ denote the set of all matrices that commute with C_γ . Any $X \in \mathfrak{X}_\gamma$ is uniquely expressed as a polynomial

$$(3.5) \quad X = \sum_{i=0}^{n-1} \alpha_i \cdot C_\gamma^i \quad (\alpha_i \in \mathbb{C})$$

in C_γ of degree at most $n-1$. This fact can be checked directly. In view of the Cayley-Hamilton theorem, the map from $\mathbb{C}[\zeta]$ sending $\varphi(\zeta)$ to $\varphi(C_\gamma)$ induces an isomorphism $\mathcal{O}_\gamma \rightarrow \mathfrak{X}_\gamma$ of affine variety. In the following, we identify $\mathcal{O}_\gamma$ with $\mathfrak{X}_\gamma$ via this map. Let Y_γ° denote the subset of $\mathbb{P}(\mathcal{O}_\gamma)$ such that the representatives $\varphi \in \mathcal{O}_\gamma - \{0\}$ satisfy the following conditions :

- (Y₀) : $\varphi(C_\gamma)$ is invertible.
- (Y₁) : $(1, n)$ component of $\varphi(C_\gamma)$ is non-zero.
- (Y₂) : $T_i(\varphi) \neq 0$ ($1 \leq i \leq n-1$).

$$(Y_3) : S_i(\varphi) \neq 0 \ (1 \leq i \leq n-1).$$

Note that (Y_1) is equivalent to the condition that $\varphi(\zeta)$ can be chosen so that it has degree $n-1$. We say φ is *normalized* if it is monic of degree $n-1$.

Remark 3.4. *It should be natural to consider the set of elements of $\mathbb{P}(\mathcal{O}_\gamma)$ satisfying (Y_1) as the centralizer of $[C_\gamma]$ in $PGL_n(\mathbb{C})$, the Langlands dual group of $SL_n(\mathbb{C})$.*

$T_i(\varphi)$ and $S_i(\varphi)$ (Definition 2.3) are given in terms of the matrix $\varphi(C_\gamma)$ as follows.

Lemma 3.5. *We have the following:*

- (1) $T_i(\varphi) = (-1)^{n-i} \xi_{1, \dots, i-1, i}^{1, \dots, i-1, n}(\varphi(C_\gamma))$.
- (2) $S_i(\varphi) = \xi_{1, \dots, i}^{1, \dots, i}(\varphi(C_\gamma))$.

Proof. Let us define $\mathbf{c} : \mathcal{O}_\gamma \rightarrow \mathbb{C}^n$ by sending polynomial to the remainder with respect to f_γ and expand it with basis $1, \zeta, \dots, \zeta^{n-1}$, in particular $\zeta^i \mapsto \mathbf{e}_{i+1}$ ($1 \leq i \leq n-1$). Then the i th row of $\varphi(C_\gamma)$ is ${}^t\mathbf{b}_{i-1}$, and $\mathbf{a}_i = \mathbf{e}_{i+1}$. Now the formulas are easily obtained. $\square$

3.4. Construction of $\alpha : Z_\gamma^\circ \rightarrow Y_\gamma^\circ$.

Proposition 3.6. *Let $(z, Q) \in Z_\gamma$ and denote $L(z, Q)$ by L .*

(1) *There is a matrix $R = (r_{ij})_{1 \leq i, j \leq n}$ in $\mathbf{B}\sigma$, unique up to scholar, such that $LR = RC_\gamma$. Moreover, for such matrix R , by multiplying some non-zero constant if needed, we have*

$$(3.6) \quad \det(R) = (-1)^{n(n-1)/2} Q_1^{n-1} Q_2^{n-2} \cdots Q_{n-1},$$

$$(3.7) \quad r_{i+1, i} = (-1)^{i-1} Q_1 \cdots Q_i \quad (1 \leq i \leq n-1).$$

(2) *There is a unique matrix U in $\mathbf{N}_-\varepsilon$ such that $LU = UC_\gamma$.*

Proof. (1) Let $\Delta_{i,j}$ denote the (i, j) -minor of $\zeta B - A$, i.e. $\Delta_{i,j} = \xi_{1, \dots, \hat{i}, \dots, \hat{j}, \dots, n}^{1, \dots, \hat{j}, \dots, n}(\zeta B - A)$. It is straightforward to show the following:

$$(3.8) \quad (-1)^{1+j} \Delta_{1,j} = (\zeta^{n-1} + \cdots + (-1)^j z_{j+1} \cdots z_n \cdot \zeta^{j-1}) \cdot \prod_{i=1}^{j-1} Q_i z_i,$$

$$(3.9) \quad \Delta_{n,j} = \zeta^{j-1} + \cdots + (-1)^{j-1} z_1 \cdots z_{j-1}.$$

Note in particular that $\Delta_{1,1}$ is monic of degree $n-1$, and $\Delta_{n,1} = 1$.

If we define a vector

$$(3.10) \quad \mathbf{v}_- := {}^t(\Delta_{1,1}, -\Delta_{1,2}, \dots, (-1)^{n-1} \Delta_{1,n})$$

in $\mathbb{C}[\zeta]^n$, then by the Laplace expansion theorem we have

$$(3.11) \quad (\zeta B - A)\mathbf{v}_- = {}^t(\det(\zeta B - A), 0, \dots, 0) = {}^t(f_\gamma(\zeta), 0, \dots, 0).$$

Let $\mathbf{w}_- := B\mathbf{v}_-$. We apply the natural projection $\mathbb{C}[\zeta] \rightarrow \mathcal{O}_\gamma$ to both hand sides of (3.11). Then we have the following equation in $\mathcal{O}_\gamma^n$:

$$(3.12) \quad (\zeta \cdot 1 - L)\mathbf{w}_- = \mathbf{0}.$$

We can write $\mathbf{w}_- = R\mathbf{v}_0$ by a unique matrix $R \in M_n(\mathbb{C})$, with $\mathbf{v}_0 = {}^t(1, \zeta, \dots, \zeta^{n-1})$. Noting that $\zeta \cdot \mathbf{v}_0 = C_\gamma \mathbf{v}_0$, (3.12) is $(RC_\gamma - LR)\mathbf{v}_0 = \mathbf{0}$, and thereby we have $RC_\gamma - LR = 0$.

We need to check that $R \in \mathbf{B}\sigma$. If we write $\mathbf{v}_- = R_0 \mathbf{v}_0$ with $R_0 \in GL_n(\mathbb{C})$ then we can see from (3.8) that R_0 is upper triangular such that n th column of R_0 is ${}^t(1, Q_1 z_1, Q_1 Q_2 z_1 z_2, \dots, Q_1 \cdots Q_{n-1} z_1 \cdots z_{n-1})$. Consequently, $R = BR_0$ has the desired form. We also have (3.7) because (j, j) entry of R_0 is $(-1)^j z_{j+1} \cdots z_n \prod_{i=1}^{j-1} Q_i z_i$.

If R' also satisfies $R'C_\gamma - LR' = 0$, then by writing $R = X\sigma, R' = X'\sigma$ with $X, X' \in \mathbf{B}$, $X^{-1}X'$ commute with $\sigma C_\gamma \sigma^{-1}$. By direct calculations, one sees that this commutativity means $X^{-1}X'$ is a scalar matrix.

Next we show (3.6). Since B is unitriangular, we have $\det(R) = \det(R_0)$. This is calculated by (3.8) as

$$\prod_{j=1}^n ((-1)^j z_{j+1} \cdots z_n \cdot \prod_{i=1}^{j-1} Q_i z_i) = (-1)^{n(n-1)/2} (z_1 \cdots z_n)^{n-1} Q_1^{n-1} Q_2^{n-2} \cdots Q_{n-1}.$$

Since $z_1 \cdots z_n = 1$, we have (3.6).

(2) We define $\mathbf{v}_+ \in \mathbb{C}[\zeta]^n$ by

$$(3.13) \quad \mathbf{v}_+ := {}^t(\Delta_{n,1}, -\Delta_{n,2}, \dots, (-1)^{n-1} \Delta_{n,n}).$$

If we set $\mathbf{w}_+ = B\mathbf{v}_+$ then we have $(\zeta \cdot 1 - L)\mathbf{w}_+ = 0$ in $\mathcal{O}_\gamma^n$. We write $\mathbf{w}_+ = U\mathbf{v}_0$. Then we have $LU = UC_\gamma$ by the same reason. From (3.9) we see that $U \in \mathbf{N}_{-\varepsilon}$. The uniqueness is straightforward. $\square$

Now we construct a morphism $\alpha^\circ : Z_\gamma^\circ \rightarrow Y_\gamma^\circ$. Let $(z, Q) \in Z_\gamma^\circ$ and $L = L(z, Q)$ be the corresponding Lax matrix (Proposition 3.3). Let U, R be matrices constructed in Proposition 3.6. Note that R is invertible because we assume (Z_4) and we have (3.6). We have $L = RC_\gamma R^{-1} = UC_\gamma U^{-1}$. So $U^{-1}R \in \mathfrak{X}_\gamma \simeq \mathcal{O}_\gamma$. Let us write $U^{-1}R = \varphi(C_\gamma)$ by a polynomial φ in ζ as (3.5). Set $\alpha^\circ(L) := [\varphi] \in \mathbb{P}(\mathcal{O}_\gamma)$. From the form of matrices R, U , one sees that the $(1, n)$ entry of $U^{-1}R$ is 1, so φ is monic of degree $n-1$. Thus (Y_1) holds for $\varphi(C_\gamma)$. We have (Y_0) since R is invertible. Since $\varphi(C_\gamma)$ is factorized as $U^{-1}R$ as in Proposition 3.2, (Y_2) holds in view of Lemma 3.5 (1). Finally we check (Y_3) . We use Lemma 3.5 (2). From similar calculation of the proof of (3.6), the i -th principle minor of $U^{-1}R (= \varphi(C_\gamma))$ is written as

$$(-1)^i z_{i+1} \cdots z_n Q_1^{i-1} Q_2^{i-2} \cdots Q_{i-1},$$

which is non-zero by assumption (Z_3) . Thus $\alpha^\circ(L)$ is an element of Y_γ° .

Proposition 3.7. *We have $\alpha|_{Z_\gamma^\circ} = \alpha^\circ$.*

Proof. Since $\varphi(C_\gamma) = U^{-1}R$, we have in $\mathcal{O}_\gamma^n$ the following:

$$\mathbf{w}_- = R\mathbf{v}_0 = U\varphi(C_\gamma)\mathbf{v}_0 = U\varphi(\zeta)\mathbf{v}_0 = \varphi(\zeta)U\mathbf{v}_0 = \varphi(\zeta)\mathbf{w}_+,$$

and thereby $\mathbf{v}_- = \varphi(\zeta)\mathbf{v}_+$. By comparing the first component of the both hand sides of this equality, we have $\Delta_{1,1} = \varphi(\zeta)$. $\square$

3.5. Construction of $\beta : Y_\gamma^\circ \rightarrow Z_\gamma^\circ$. Let φ be a normalized element in Y_γ° . Since we assume (Y_2) , from Proposition 3.2, we have unique R, U such that

$$(3.14) \quad \varphi(C_\gamma) = U^{-1}R.$$

From (Y_0) , we see that R is invertible. Now we have $UC_\gamma U^{-1} = RC_\gamma R^{-1} =: L$. Set $\beta([\varphi]) := L$. We claim that L is an element of Z° . It is easy to check that L satisfies (Z_1) and (Z_2) by using $L = UC_\gamma U^{-1}$ and $L = RC_\gamma R^{-1}$ respectively. (Z_3) follows from (Y_3) by the following.

Lemma 3.8. *Let φ and L as above. Then we have*

$$(3.15) \quad \xi_{i+1,\dots,n}^{i+1,\dots,n}(L) = -\frac{S_i(\varphi)}{T_i(\varphi)} \quad (1 \leq i \leq n-1).$$

Proof. Note that for any element R in $\mathbf{B}\sigma$, we have

$$\xi_{i+1,\dots,n}^{i+1,\dots,n}(RC_\gamma R^{-1}) = (-1)^{n-1} \frac{\xi_{1,\dots,i}^{1,\dots,i}(R)}{r_{21}r_{32} \cdots r_{i,i-1}}.$$

Now using (3.2) we have $r_{21}r_{32} \cdots r_{i,i-1} = (-1)^{n-i+i(i+1)/2} T_i(\varphi)$, and

$$\xi_{1,\dots,i}^{1,\dots,i}(R) = (-1)^{i(i-1)/2} \xi_{1,\dots,i}^{1,\dots,i}(U^{-1}R).$$

Thus we have the result from Lemma 3.5 (2). $\square$

Finally we need to show (Z_4) . It is assured by the fact that R is invertible. Let $L = L(z, Q)$. Note that $\det(R) = (-1)^{n-1} r_{2,1} r_{3,2} \cdots r_{n,n-1} \neq 0$. Note also that $(i+1, i)$ entry of L^{-1} is $-Q_i$ (cf. (3.4)), which is the ratio $r_{i+1,i}/r_{i,i-1}$ with $r_{1,0} = -1$, and thus non-zero. Thus L is in Z° . Since L is conjugate to C_γ we have $\beta([\varphi]) = L \in Z_\gamma^\circ$.

Proposition 3.9. *The morphisms α° and β are inverse to each other.*

Proof. Let $L \in Z_\gamma^\circ$. We take U, R such that $L = UC_\gamma U^{-1} = RC_\gamma R^{-1}$. Then $\varphi(\zeta) = \Delta_{1,1}$ satisfies $\varphi(C_\gamma) = U^{-1}R$. Then $\alpha(L) = [\varphi]$ and $\beta([\varphi]) = UC_\gamma U^{-1} = L$. On the other hand, let $[\varphi] \in Y_\gamma^\circ$. We assume φ is normalized. We have $\varphi(C_\gamma) = U^{-1}R$. Then $\beta([\varphi]) = UC_\gamma U^{-1} = RC_\gamma R^{-1} =: L$. Then $\alpha^\circ(L)$ is given by $U^{-1}R$. $\square$

3.6. Explicit form of Φ_n .

Proposition 3.10. *Let $\Phi_n : \mathbb{C}[Z_\gamma^\circ] \rightarrow \mathbb{C}[Y_\gamma^\circ]$ be the associated isomorphism of coordinate rings. Then we have*

$$(3.16) \quad \Phi_n(z_i) = \frac{T_i(\varphi)S_{i-1}(\varphi)}{S_i(\varphi)T_{i-1}(\varphi)},$$

$$(3.17) \quad \Phi_n(Q_i) = \frac{T_{i-1}(\varphi)T_{i+1}(\varphi)}{T_i^2(\varphi)}.$$

Proof. It is easy to show $\xi_{i,\dots,n}^{i,\dots,n}(L) = z_i \cdots z_n$. Then (3.16) follows from (3.15) and Lemma 3.5. To prove (3.17) we use (3.2), (3.7) to have

$$Q_i = -\frac{r_{i+1,i}}{r_{i,i-1}} = \frac{\xi_{1,\dots,i-1,n}^{1,\dots,i-1,n}(X)}{\xi_{1,\dots,i-2,n}^{1,\dots,i-2,n}(X)} \cdot \frac{\xi_{1,\dots,i-3,n}^{1,\dots,i-3,n}(X)}{\xi_{1,\dots,i-2,i-1}^{1,\dots,i-2,n}(X)}.$$

with $X = \varphi(C_\gamma)$. Then from Lemma 3.5 (1), we have (3.17). $\square$

Remark 3.11. *We observe that each component of L expressed as a rational function of $T_i(\varphi)$'s and $S_i(\varphi)$'s has no pole along the divisor defined by $S_i(\varphi) = 0$. This is mysterious because z_i has pole along the divisor. We do not know any explanation of this phenomena.*

4. DUAL STABLE GROTHENDIECK POLYNOMIALS AS τ -FUNCTIONS

In this section we prove (2.5). It shows that the functions T_d, S_d defined as the determinants are, in the unipotent case, written in terms of the dual Grothendieck polynomials associated to the rectangle R_d .

4.1. The determinant formula for dual stable Grothendieck polynomials.

Let $U_0 \subset \mathbb{P}(\mathcal{O}_{\text{uni}})$ be the affine subspace defined by $c_0 \neq 0$. We identify the coordinate ring $\mathbb{C}[U_0]$ with the ring $\Lambda_{(n)} = \mathbb{C}[h_1, \dots, h_{n-1}]$ through the identification $c_i/c_0 \leftrightarrow h_i$ (see (2.3) for the definition of c_i).

We fix d such that $1 \leq d \leq n$. For $f_1, f_2, \dots, f_d \in \mathcal{O}_{\text{uni}}$, define $[f_1, f_2, \dots, f_d] \in \mathbb{C}[U_0] = \Lambda_{(n)}$ by the formula

$$\begin{aligned} & [f_1, f_2, \dots, f_d](\varphi) \\ &= c_0^{-d} \cdot |\mathbf{c}(f_1\varphi), \mathbf{c}(f_2\varphi), \dots, \mathbf{c}(f_d\varphi), \mathbf{c}(1), \mathbf{c}(\zeta), \dots, \mathbf{c}(\zeta^{n-d-1})| \quad (\varphi \in U_0). \end{aligned}$$

If $f_j\varphi \in U_0$ is expressed as $f_j\varphi = \sum_{i=0}^{n-1} a_{i,j}(\zeta-1)^i \pmod{(\zeta-1)^n}$, we have

$$(4.18) \quad [f_1, \dots, f_d](\varphi) = (-1)^{d(n-d)} c_0^{-d} \cdot \det(a_{n-d+i-1,j})_{i,j=1}^d.$$

Note that for $m \geq 0$, ζ^{-m} makes sense as an element of $\mathcal{O}_{\text{uni}}$ since $(1-\zeta)^n = 0$. We have $\zeta^{-m} = \sum_{l=0}^{n-1} (-1)^l \binom{m+l-1}{l} (\zeta-1)^l$.

Lemma 4.1. *For a partition $\lambda \subset R_d$, we have*

$$(4.19) \quad [(1-\zeta)^{n-1-\lambda_d} \zeta^{-d+1}, (1-\zeta)^{n-2-\lambda_{d-1}} \zeta^{-d+2}, \dots, (1-\zeta)^{n-d-\lambda_1}] = g_\lambda.$$

Proof. For $a, b \geq 0$, we write $(1-\zeta)^a \zeta^{-b} \varphi = \sum_{i=0}^{n-1} c_i^{a,b} (\zeta-1)^i \pmod{(\zeta-1)^n}$. Then we have

$$c_i^{a,b} = \sum_{m=0}^{\infty} (-1)^{i+m} \binom{-b}{m} h_{i-a-m},$$

where we understand $h_m = 0$ for $m < 0$. By using (4.18), the left hand side of (4.19) can be calculated as follows:

$$\begin{aligned} & (-1)^{d(n-d)} \det \left(c_{n-d+j-1}^{n-i-\lambda_{d-i+1}, d-i} \right)_{i,j=1}^d \\ &= (-1)^{d(n-d)} \det \left(\sum_{m=0}^{\infty} (-1)^{n-d+j-1+m} \binom{i-d}{m} h_{\lambda_{d-i+1}+j-(d-i+1)-m} \right)_{i,j=1}^d \\ &= (-1)^{d(n-d)+\sum_{j=1}^d (n-d+j-1)+d(d-1)/2} \det \left(\sum_{m=0}^{\infty} (-1)^m \binom{1-i}{m} h_{\lambda_i+j-i-m} \right)_{i,j=1}^d \\ &= g_\lambda. \end{aligned}$$

□

Proposition 4.2. *We have $\tau_d = g_{R_d}$.*

Proof. Note that $M_\zeta \mathbf{c}(f) = \mathbf{c}(f\zeta)$ with $M_\zeta := E_n + J$. We have

$$\begin{aligned} \tau_d &= c_0^{-d} \cdot |\mathbf{c}(\varphi), \dots, \mathbf{c}(\varphi\zeta^{d-1}), \mathbf{c}(\zeta^{d-1}), \dots, \mathbf{c}(\zeta^{n-2})| \\ &= c_0^{-d} \cdot |M_\zeta|^{d-1} \left| \mathbf{c}(\varphi\zeta^{-(d-1)}), \dots, \mathbf{c}(\varphi), \mathbf{c}(1), \dots, \mathbf{c}(\zeta^{n-d-1}) \right| \\ &= [\zeta^{-(d-1)}, \zeta^{-(d-2)}, \dots, 1] \\ &= [(1-\zeta)^{d-1} \zeta^{-(d-1)}, (1-\zeta)^{d-2} \zeta^{-(d-2)}, \dots, 1]. \end{aligned}$$

The last equality follows from the fact that $(1-\zeta)^m \zeta^{-m} - \zeta^{-m}$ ($m \geq 1$) is a linear combination of $1, \zeta^{-1}, \dots, \zeta^{-m+1}$. Then from Lemma 4.6 we obtain $\tau_d = g_{R_d}$. □

4.2. A useful family of determinants. Here we introduce some useful family of determinant formulas for symmetric polynomials which is suitable for our representation of K -theoretic Peterson isomorphism.

Definition 4.3. For d -tuple $(\theta_1, \dots, \theta_d)$ of integers and d -tuple $(a_1, \dots, a_d)$ of non-negative integers, define

$$D \begin{pmatrix} \theta_1, \theta_2, \dots, \theta_d \\ a_1, a_2, \dots, a_d \end{pmatrix} := [(1 - \zeta)^{a_1} \zeta^{-\theta_1}, (1 - \zeta)^{a_2} \zeta^{-\theta_2}, \dots, (1 - \zeta)^{a_d} \zeta^{-\theta_d}].$$

We also denote $D \begin{pmatrix} \theta_1, \theta_2, \dots, \theta_d \\ 0, 0, \dots, 0 \end{pmatrix}$ simply by $D(\theta_1, \theta_2, \dots, \theta_d)$.

Lemma 4.4. $D \begin{pmatrix} \theta_1, \theta_2, \dots, \theta_d \\ a_1, a_2, \dots, a_d \end{pmatrix} \in \Lambda_{(n)}$ is uniquely determined by the following recursive formulas:

- (1) $D \begin{pmatrix} \dots, \theta_i, \dots, \theta_j, \dots \\ \dots, a_i, \dots, a_j, \dots \end{pmatrix} = -D \begin{pmatrix} \dots, \theta_j, \dots, \theta_i, \dots \\ \dots, a_j, \dots, a_i, \dots \end{pmatrix},$
- (2) $D \begin{pmatrix} \dots, \theta_i, \dots \\ \dots, a_i, \dots \end{pmatrix} = D \begin{pmatrix} \dots, \theta_i - 1, \dots \\ \dots, a_i, \dots \end{pmatrix} + D \begin{pmatrix} \dots, \theta_i, \dots \\ \dots, a_i + 1, \dots \end{pmatrix},$
- (3) $D \begin{pmatrix} \dots, \theta, \dots \\ \dots, n, \dots \end{pmatrix} = 0,$
- (4) $D \begin{pmatrix} 0, \dots, 0 \\ n - \lambda_d - 1, n - \lambda_{d-1} - 2, \dots, n - \lambda_1 - d \end{pmatrix} = s_\lambda, \text{ where } s_\lambda \text{ is the Schur function.}$

We have useful formulas such as

$$(4.20) \quad D \begin{pmatrix} d-1, d-2, \dots, 0 \\ 0, 0, \dots, 0 \end{pmatrix} = D \begin{pmatrix} d-1, d-2, \dots, 0 \\ d-1, d-2, \dots, 0 \end{pmatrix} = g_{R_d},$$

$$(4.21) \quad D \begin{pmatrix} d-1, \dots, d-2, \dots, 0 \\ n - \lambda_d - 1, n - \lambda_{d-1} - 2, \dots, n - d - \lambda_1 \end{pmatrix} = g_\lambda, \quad (\lambda \subset R_d)$$

$$(4.22) \quad D \begin{pmatrix} \dots, \theta, \dots \\ \dots, a, \dots \end{pmatrix} = \sum_i (-1)^i \binom{p}{i} D \begin{pmatrix} \dots, \theta + p, \dots \\ \dots, a + i, \dots \end{pmatrix}.$$

4.3. Lattice paths method. Now we prove the following.

Proposition 4.5. We have $\sigma_d = \sum_{\mu \subset R_d} g_\mu$.

Since we have Lemma 4.1, it suffices to show the following.

Lemma 4.6. We have

$$(4.23) \quad D \begin{pmatrix} d, d-1, \dots, 1 \\ d-1, d-2, \dots, 0 \end{pmatrix} = \sum_{n > a_1 > a_2 > \dots > a_d \geq 0} D \begin{pmatrix} d-1, d-2, \dots, 0 \\ a_1, a_2, \dots, a_d \end{pmatrix}.$$

Proof. Let

$$K_{a,i}^\theta := \binom{i - a + \theta - 1}{\theta - 1} = \left(\begin{array}{c} \text{The number of weakly increasing sequences} \\ a \leq x_1 \leq x_2 \leq \dots \leq x_\theta \leq i \end{array} \right).$$

By using (4.22) repeatedly, we have

$$\begin{aligned} D \begin{pmatrix} \theta_1, \dots, \theta_d \\ a_1, \dots, a_d \end{pmatrix} &= \sum_{n > i_1 > \dots > i_d \geq 0} \sum_{\sigma \in S_d} \text{sgn}(\sigma) \prod_{j=1}^d K_{a_j, i_{\sigma(j)}}^{\theta_j} D \begin{pmatrix} 0, \dots, 0 \\ i_1, \dots, i_d \end{pmatrix} \\ &= \sum_{n > i_1 > \dots > i_d \geq 0} \det(K_{a_l, i_m}^{\theta_l})_{1 \leq l, m \leq d} \cdot D \begin{pmatrix} 0, \dots, 0 \\ i_1, \dots, i_d \end{pmatrix}. \end{aligned}$$

Especially,

$$\begin{aligned} D \begin{pmatrix} d, d-1, \dots, 1 \\ d-1, d-2, \dots, 0 \end{pmatrix} &= \sum_{n > i_1 > \dots > i_d \geq 0} \det(K_{d-l, i_m}^{d-l+1})_{1 \leq l, m \leq d} \cdot D \begin{pmatrix} 0, \dots, 0 \\ i_1, \dots, i_d \end{pmatrix}, \\ D \begin{pmatrix} d-1, d-2, \dots, 0 \\ a_1, a_2, \dots, a_d \end{pmatrix} &= \sum_{n > i_1 > \dots > i_d \geq 0} \det(K_{a_l, i_m}^{d-l})_{1 \leq l, m \leq d} \cdot D \begin{pmatrix} 0, \dots, 0 \\ i_1, \dots, i_d \end{pmatrix}. \end{aligned}$$

Hence, it suffices to prove

$$(4.24) \quad \det(K_{d-l, i_m}^{d-l+1})_{l, m} = \sum_{n > a_1 > \dots > a_d \geq 0} \det(K_{a_l, i_m}^{d-l})_{l, m}$$

for arbitrary $i_1, \dots, i_d$.

Now consider the plane lattice in Figure 1. Let A_j ($j = 1, \dots, d$) be the point with coordinates $(d-j+1, d-j)$ and B_j be the point with $(0, i_j)$. We immediately find that the number of shortest paths from A_l to B_m on the lattice is K_{d-j, i_m}^{d-j+1} . By the Lindström-Gessel-Viennot lemma [11], the determinant $\det(K_{d-l, i_m}^{d-l+1})_{l, m}$ equals to the number of shortest non-intersecting paths from the source set $\{A_1, \dots, A_d\}$ to the target set $\{B_1, \dots, B_d\}$. (See Figure 1).




FIGURE 1. ($d = 5$). An example of non-intersecting paths from $\{A_1, \dots, A_5\}$ to $\{B_1, \dots, B_5\}$. The plane lattice lacks the line on the 0-th column. Here, $(i_1, i_2, i_3, i_4, i_5) = (9, 8, 5, 4, 2)$. It corresponds to non-intersecting paths from $\{C_1, \dots, C_5\}$ to $\{B_1, \dots, B_5\}$ with $(a_1, a_2, a_3, a_4, a_5) = (7, 6, 5, 3, 2)$.

Let C_j ($j = 1, \dots, d$) be the point with coordinates $(d-j, a_j)$. Similarly as above, the determinant $\det(K_{a_j, i_m}^{d-j})$ equals to the number of shortest non-intersecting paths from the source set $\{C_1, \dots, C_d\}$ to the target set $\{B_1, \dots, B_d\}$. (See Figure 1).

Denote

$\mathfrak{X} = (\text{the set of shortest non-intersecting paths from } \{A_1, \dots, A_d\} \text{ to } \{B_1, \dots, B_d\}),$

$$\mathfrak{Y}^{(a_1, \dots, a_d)} = \left(\begin{array}{c} \text{The number of shortest non-intersecting paths} \\ \text{from } \{C_1, \dots, C_d\} \text{ to } \{B_1, \dots, B_d\} \text{ with } C_j = (d-j, a_j) \end{array} \right).$$

There exists a natural one-to-one correspondence

$$\bigcup_{n > a_1 > \dots > a_d \geq 0} \mathfrak{Y}^{(a_1, \dots, a_d)} \rightarrow \mathfrak{X}$$

which associates

$$(\mathcal{P}_1, \dots, \mathcal{P}_d) \in \mathfrak{Y}^{(a_1, \dots, a_d)}, \quad (\mathcal{P}_j \text{ is a path from } C_j \text{ to } B_j)$$

with the set of non-intersecting paths $(\mathcal{Q}_1, \dots, \mathcal{Q}_d) \in \mathfrak{X}$, where $\mathcal{Q}_j$ is the path which is obtained by adding to $\mathcal{P}_j$ a vertical segment from A_j to $(d-j+1, a_j)$ and a horizontal edge from $(d-j+1, a_j)$ to $(d-j, a_j)$ (see Figure 1). Counting the cardinalities of these sets concludes (4.24). $\square$

5. QUANTUM GROTHENDIECK POLYNOMIALS OF GRASSMANNIAN TYPE

We fix an integer d with $1 \leq d \leq n$. Let $Gr_d(\mathbb{C}^n)$ denote the Grassmannian of d -dimensional subspaces of $\mathbb{C}^n$. We will show how a quantization map for the Grassmannian $Gr_d(\mathbb{C}^n)$ can be interpreted in our context. Theorem 5.6 is the main result whose proof is given in the next section. As an application, we obtain Theorem 1.6.

5.1. A result on K -theoretic Littlewood-Richardson rule. For partitions λ, μ, ν , the K -theoretic Littlewood-Richardson coefficients is the integer $c_{\lambda, \mu}^\nu$ defined by

$$G_\lambda \cdot G_\mu = \sum_{\nu} (-1)^{|\nu| - |\lambda| - |\mu|} c_{\lambda, \mu}^\nu \cdot G_\nu,$$

where ν runs for all partitions. Buch's formula [5] gives $c_{\lambda, \mu}^\nu$ as the number of set-valued tableaux which satisfy a certain property. Here we explain a rule of Ikeda-Shimazaki [17], which is equivalent to Buch's rule. Let T be a set-valued tableau. The *column word* $cw(T)$ of T is obtained by reading each column from top to bottom starting from the right most column to the left, where letters in the set filled in a box are read in the decreasing order. A word *builds* ν on λ if ν is constructed by adding a box in the row associated to each entry of the word one by one while keeping the shape being a partition. The coefficient $c_{\lambda, \mu}^\nu$ is the number of set-valued tableaux T of shape μ such that $cw(T)$ builds ν from λ (in [17], such T is called λ -good tableaux of shape μ of content $\nu - \lambda$).

The following is stated without proof in [6] (proof of Corollary 1). We include a proof for completeness.

Proposition 5.1. *Let λ be a partition such that $\lambda \subset R_d$. For any partition μ , we have*

$$(5.1) \quad c_{\lambda, \mu}^{R_d} = \delta_{\lambda^\vee, \mu}.$$

Proof. (T. Matsumura) Let T be a set-valued tableau such that $cw(T)$ builds R_d on λ . We prove that T is the ordinary semistandard tableau T_0 such that i th column of T_0 is filled with $d, d-1, \dots, r_i+1$ from the bottom to top, where r_i is the number

There exist a unique family $\{\mathfrak{G}_w(x) \mid w \in S_n\}$ of polynomials such that

$$\pi_i \mathfrak{G}_w = \begin{cases} \mathfrak{G}_{ws_i} & \text{if } \ell(ws_i) = \ell(w) - 1 \\ \mathfrak{G}_w & \text{if } \ell(ws_i) = \ell(w) + 1. \end{cases}$$

5.3. Quantization map of $K(Gr_d(\mathbb{C}^n))$ and K -Peterson isomorphism. Let $\hat{\Lambda}$ denote the $\mathbb{C}$ -span of the stable Grothendieck polynomials. This is a completion of the ring Λ . It was shown in [5] that $\hat{\Lambda}$, denoted as Γ in the paper, is closed under multiplication. Denote by J_d the ideal of $\hat{\Lambda}$ defined as

$$J_d := \{f \in \hat{\Lambda} \mid f^\perp \cdot g_{R_d} = 0\}.$$

Proposition 5.3. *There is a canonical isomorphism $K(Gr_d(\mathbb{C}^n)) \simeq \hat{\Lambda}/J_d$.*

Proof. The linear span of G_μ 's such that $\mu \not\leq R_d$ is an ideal of $\hat{\Lambda}$ and $\hat{\Lambda}/I_d$ is isomorphic $K(Gr_d(\mathbb{C}^n))$ (Buch [5], Theorem 8.1). From Proposition 5.1, we see that J_d coincides with I_d . $\square$

Via the isomorphism in Proposition 5.3, we define the $\mathbb{C}$ -linear map

$$\widehat{Q}_d : K(Gr_d(\mathbb{C}^n)) \rightarrow \mathcal{QK}(Fl_n), \quad f \bmod J_d \mapsto \Phi_n^{-1}((f^\perp \cdot g_{R_d})/g_{R_d}).$$

5.4. Quantization map of Lenart–Maeno. For $1 \leq m \leq n$, define a polynomial $F_i^{(m)} \in \mathbb{C}[x, Q]$ (cf. [38, §3]) by

$$F_i^{(m)} = \sum_{\substack{I \subset \{1, \dots, m\} \\ \#I=i}} \prod_{j \in I} (1 - x_j) \prod_{j \in I, j+1 \notin I} (1 - Q_j),$$

where $Q_n := 0$. Note that $F_i^{(n)}$ is nothing but F_i with $z_j = 1 - x_j$.

In [38], Lenart and Maeno introduced the *quantization map* $\widehat{Q}$ and defined the *quantum Grothendieck polynomials* by using it. Let e_i be the i th elementary symmetric polynomial. Let $f_i^{(j)} = e_i(1 - x_1, \dots, 1 - x_j)$ for $1 \leq i, j \leq n$. The following presentation is well-known (recall that x_i is the K -theoretic first Chern class $c_1(\mathcal{L}_i^\vee) := 1 - [\mathcal{L}_i]$ of the dual of the tautological line bundle $\mathcal{L}_i$):

$$(5.2) \quad K(Fl_n) \simeq \mathbb{C}[x_1, \dots, x_n] / \langle e_i(x_1, \dots, x_n) \mid 1 \leq i \leq n \rangle.$$

Note that the ideal $\langle e_i(x_1, \dots, x_n) \mid 1 \leq i \leq n \rangle$ is also generated by $f_i^{(n)} - \binom{n}{i}$ ($1 \leq i \leq n$). Let L_n be the $\mathbb{C}$ -vector subspace of $\mathbb{C}[x_1, \dots, x_n]$ generated by the elements

$$(5.3) \quad f_{i_1}^{(1)} f_{i_2}^{(2)} \dots f_{i_{n-1}}^{(n-1)} \quad (0 \leq i_j \leq j).$$

There exists a canonical isomorphism $K(Fl_n) \simeq L_n$ of $\mathbb{C}$ -vector spaces ([38]).

Definition 5.4 ([38]). *The quantization map $\widehat{Q} : K(Fl_n) \rightarrow \mathcal{QK}(Fl_n)$ is the $\mathbb{C}$ -linear map defined by*

$$(5.4) \quad \widehat{Q}(f_{i_1}^{(1)} f_{i_2}^{(2)} \dots f_{i_{n-1}}^{(n-1)}) := F_{i_1}^{(1)} F_{i_2}^{(2)} \dots F_{i_{n-1}}^{(n-1)} \quad (0 \leq i_j \leq j).$$

Remark 5.5. *The definition of $\widehat{Q}$ given in [38] is equivalent to Definition 5.4 (see [38, Proposition 3.16]).*

The quantum Grothendieck polynomial $\mathfrak{G}_w^Q$, for $w \in S_n$, is defined as

$$\mathfrak{G}_w^Q = \widehat{Q}(\mathfrak{G}_w).$$

For $f \in \hat{\Lambda}$, let $f(x_1, \dots, x_d)$ denotes the polynomial by setting $x_i = 0$ for $i > d$ in the symmetric function $f \in \hat{\Lambda}$. Let $\pi : Fl_n \rightarrow Gr_d(\mathbb{C}^n)$ be the projection sending $V_\bullet$ to V_d . The induced morphism $\pi^* : K(Gr_d(\mathbb{C}^n)) \simeq \hat{\Lambda}/J_d \hookrightarrow K(Fl_n)$ is given by $f \bmod J_d \mapsto f(x_1, \dots, x_d)$.

The main statement of this section is the following. The proof is given in §6.

Theorem 5.6. *The following diagram commutes*

$$\begin{array}{ccc} \hat{\Lambda}/J_d \simeq K(Gr_d(\mathbb{C}^n)) & \xrightarrow{\pi^*} & K(Fl_n) \\ & \searrow \widehat{Q}_d \quad \swarrow \widehat{Q} & \\ & \mathcal{QK}(Fl_n) & \end{array}$$

Corollary 5.7. *Let $\lambda \subset R_d$. We have $\widehat{Q}_d(G_\lambda \bmod J_d) = \mathfrak{G}_{w_{\lambda,d}}^Q$.*

Proof. It is known that $G_\lambda(x_1, \dots, x_d) = \mathfrak{G}_{\lambda,d}$ (Buch [5], §8), so we have

$$\widehat{Q}_d(G_\lambda \bmod J_d) = \widehat{Q}(G_\lambda(x_1, \dots, x_d)) = \widehat{Q}(\mathfrak{G}_{\lambda,d}) = \mathfrak{G}_{\lambda,d}^Q.$$

□

Now we prove Theorem 1.6.

Proof. Corollary 5.7 is equivalent to

$$\Phi_n(\mathfrak{G}_{w_{\lambda,d}}^Q) = \frac{G_\lambda^\perp \cdot g_{R_d}}{g_{R_d}}.$$

So we need to show $G_\lambda^\perp \cdot g_{R_d} = g_{\lambda^\vee}$, which is equivalent to Proposition 5.1. □

6. PROOF OF THEOREM 5.6

6.1. Outline of the proof. Let λ be partition such that $\lambda \subset R_d$. We define a quantized version of Schur polynomial

$$S_{\lambda,d}^Q := \det(F_{\lambda'_i - i + j}^{(d+j-1)})_{i,j=1}^{\ell(\lambda')}.$$

Note that the polynomial $S_{\lambda,d}^Q$ is an element in $\mathbb{Z}[Q][x_1, \dots, x_d]^{S_d}$.

Proposition 6.1. *We have*

$$(6.1) \quad \widehat{Q}(s_\lambda(1 - x_1, \dots, 1 - x_d)) = S_{\lambda,d}^Q.$$

Proof. Let $e_i^{(j)} = e_i(z_1, \dots, z_j)$. We know the dual Jacobi-Trudi formula

$$s_\lambda(z_1, \dots, z_d) = \det(e_{\lambda'_i - i + j}^{(d)})_{i,j=1}^{\ell(\lambda')}.$$

Since $e_i^{(j)} = e_i^{(j-1)} + z_j e_{i-1}^{(j-1)}$, it is easy to show $s_\lambda(z_1, \dots, z_d) = \det(e_{\lambda'_i - i + j}^{(d+j-1)})_{i,j=1}^{\ell(\lambda')}$. Now by substituting $z_i = 1 - x_i$ to this equality, we have

$$(6.2) \quad s_\lambda(1 - x_1, \dots, 1 - x_d) = \det(f_{\lambda'_i - i + j}^{(d+j-1)})_{i,j=1}^{\ell(\lambda')}.$$

One observes that each term of the expansion on the right hand side of (6.2) is of the form (5.3). Then (6.1) follows from the definition of $\widehat{Q}$. □

Let $p_i \in \Lambda$ be the i th power sum symmetric function ([40]). We consider the ring homomorphism $\kappa_d : \Lambda \rightarrow \Lambda$ given by

$$(6.3) \quad \kappa_d(p_i) := d - \binom{i}{1}p_1 + \binom{i}{2}p_2 - \cdots + (-1)^i \binom{i}{i}p_i.$$

Recall that each element of Λ is a symmetric formal power series in $x = (x_1, x_2, \dots)$. κ_d is given by $x_i \mapsto 1 - x_i$ ($1 \leq i \leq d$), $x_j \mapsto x_j$ ($j > d$). Thus obviously κ_d is an involution. We will show the following in the sequel of this section.

Proposition 6.2. *We have*

$$(6.4) \quad \Phi_n(S_{\lambda,d}^Q) = \frac{\kappa_d(s_\lambda)^\perp \cdot g_{R_d}}{g_{R_d}}.$$

The equation (6.4) is equivalent to

$$(6.5) \quad \widehat{Q}_d(\kappa_d(s_\lambda) \bmod J_d) = S_{\lambda,d}^Q.$$

On the other hand, the element $\kappa_d(s_\lambda) \bmod J_d$ of $\hat{\Lambda}/J_d$ is mapped to $s_\lambda(1 - x_1, \dots, 1 - x_d) \in L_n \simeq K(Fl_n)$. Thus we have

$$\widehat{Q}(\pi^*(\kappa_d(s_\lambda) \bmod J_d)) = \widehat{Q}(s_\lambda(1 - x_1, \dots, 1 - x_d)) = S_{\lambda,d}^Q.$$

Since $\kappa_d(s_\lambda) \bmod J_d$ ($\lambda \subset R_d$) form a basis of $\hat{\Lambda}/J_d$, Theorem 5.6 holds.

6.2. $\Phi_n(S_{\lambda,d}^Q)$ as a ratio of determinants. In this subsection we prove the following proposition.

Proposition 6.3. *Let λ be a partition contained in R_d . We define the increasing sequence $(i_1, \dots, i_d)$ by setting*

$$(6.6) \quad i_a = \lambda_{d+1-a} + a \quad (a = 1, \dots, d).$$

Then we have

$$(6.7) \quad \Phi_n(S_{\lambda,d}^Q) = \frac{D(d - i_1, d - i_2, \dots, d - i_d)}{D(d - 1, d - 2, \dots, 0)}.$$

Remark 6.4. *Almost all the statements and proofs of this subsection make sense for arbitrary values of γ , which will be relevant when we discuss equivariant case (cf. [29]).*

Lemma 6.5. *Let $L \in Z_{\text{uni}}^\circ$, and U be the matrix constructed in Proposition 3.6. The components u_{ij} ($i > j$) of U is equal to $(-1)^{j-1} F_{i-j}^{(i-1)}$.*

Proof. We have $UC_{\text{uni}}U^{-1} = L$. Let us compare both hand sides of principal minors of $\zeta \cdot 1 - UC_{\text{uni}}U^{-1} = \zeta \cdot 1 - L$. In view of the facts that L satisfies (Z_1) , and $U \in \mathbf{N}_{-\varepsilon}$, one can show that

$$\xi_{1,\dots,i}^{1,\dots,i}(\zeta \cdot 1 - L) = \zeta^i + (-1)^i \sum_{j=1}^i u_{i+1,i-j+1} \zeta^{i-j} \quad (1 \leq i \leq n-1),$$

On the other hand, we have $\xi_{1,\dots,i}^{1,\dots,i}(\zeta \cdot 1 - L) = \zeta^i + \sum_{j=1}^i (-1)^j F_j^{(i)} \zeta^{i-j}$. Thus the lemma is proved. $\square$

Lemma 6.6 ([44]). *Let λ and $(i_1, \dots, i_d)$ be as Proposition 6.3. Let $s = \ell(\lambda')$, where λ' is the conjugate partition of λ . We define the increasing sequence $(j_1, \dots, j_s)$ by the condition*

$$\{i_1, \dots, i_d\} \cup \{j_1, \dots, j_s\} = \{1, 2, \dots, d + s\}.$$

Suppose a matrix X is decomposed as $X = Y \cdot N^{-1}$, $Y \in \mathbf{B}_-$, $N \in \mathbf{N}$. Then we have the expression

$$(6.8) \quad \xi_{j_1, \dots, j_s}^{d+1, \dots, d+s}(N) = (-1)^{|\lambda|} \cdot \frac{\xi_{1, \dots, d}^{i_1, \dots, i_d}(X)}{\xi_{1, \dots, d}^{1, \dots, d}(X)}.$$

Proof. See [44], Theorem 1.1 and its proof. $\square$

Now we prove Proposition 6.3.

Proof. Let $\{j_1, \dots, j_s\}$ as in Lemma 6.6. From Lemma 6.5, we see that

$$(6.9) \quad S_{\lambda, d}^Q = \xi_{d+1, \dots, d+s}^{j_1, \dots, j_s}(U\varepsilon).$$

We apply Lemma 6.6 as follows. Recall that we have the decomposition $\varphi(C_{\text{uni}}) = U^{-1}R$ with $R \in \mathbf{B}\sigma$ and $U \in \mathbf{N}_{-\varepsilon}$. Let $Y = {}^t(R\sigma^{-1})$, $N = {}^t(U\varepsilon)$, $X = \sigma \cdot {}^t\varphi(C_{\text{uni}})\varepsilon$. If we choose $\mathbf{c} : \mathcal{O}_{\text{uni}} \rightarrow \mathbb{C}^n$ as $\sum_{i=0}^{n-1} \alpha_i \zeta^i \pmod{(\zeta-1)^n} \mapsto {}^t(\alpha_{n-1}, \alpha_0, \dots, \alpha_{n-2})$, then we have

$$\sigma \cdot {}^t\varphi(C_{\text{uni}}) = (\mathbf{b}_0, \mathbf{b}_1, \dots, \mathbf{b}_{n-1}).$$

Now we have

$$(6.10) \quad \begin{aligned} \xi_{1, \dots, d}^{i_1, \dots, i_d}(X) &= (-1)^{\sum_{a=1}^d (i_a - 1)} |\mathbf{b}_{i_1-1}, \dots, \mathbf{b}_{i_d-1}, \mathbf{a}_{d-1}, \dots, \mathbf{a}_{n-2}| \\ &= (-1)^{\sum_{a=1}^d (i_a - 1)} |\mathbf{c}(\zeta^{i_1-1}\varphi), \dots, \mathbf{c}(\zeta^{i_d-1}\varphi), \mathbf{c}(\zeta^{d-1}), \dots, \mathbf{c}(\zeta^{n-2})| \\ &= (-1)^{\sum_{a=1}^d (i_a - 1)} |\mathbf{c}(\zeta^{i_1-d}\varphi), \dots, \mathbf{c}(\zeta^{i_d-d}\varphi), \mathbf{c}(1), \dots, \mathbf{c}(\zeta^{n-d-1})| \\ &= (-1)^{|\lambda| + d(d-1)/2} D(d - i_1, \dots, d - i_d) c_0^d, \end{aligned}$$

where we used $\sum_{a=1}^d i_a = |\lambda| + d(d+1)/2$. Since we have

$$\xi_{d+1, \dots, d+s}^{j_1, \dots, j_s}(U\varepsilon) = \xi_{j_1, \dots, j_s}^{d+1, \dots, d+s}(N),$$

formula (6.7) follows from (6.8), (6.9), and (6.10). $\square$

Remark 6.7. *The d -Grassmannian permutation $w_{\lambda, d} \in S_n$ is given by $w_{\lambda, d}(a) = i_a$ ($1 \leq a \leq d$) and $w_{\lambda, d}(d+a) = j_a$ ($1 \leq a \leq s$), and $w_{\lambda, d}(a) = a$ ($a > d+s$).*

6.3. Calculation of $\kappa_d(s_\lambda)^\perp \cdot g_{R_d}$. In view of Proposition 6.3, Proposition 6.2 is reduced to the following.

Proposition 6.8. *We have*

$$(6.11) \quad \kappa_d(s_\lambda)^\perp \cdot g_{R_d} = D(d - i_1, d - i_2, \dots, d - i_d).$$

In the rest of the paper, we will concentrate on the proof of Proposition 6.8.

6.3.1. Actions of $\kappa_d(p_i)^\perp$.

Lemma 6.9. *We have*

$$(6.12) \quad \kappa_d(p_i)^\perp \cdot D \begin{pmatrix} \theta_1, \dots, \theta_d \\ a_1, \dots, a_d \end{pmatrix} = \sum_{j=1}^d D \begin{pmatrix} \theta_1, \dots, \theta_j - i, \dots, \theta_d \\ a_1, \dots, a_j, \dots, a_d \end{pmatrix}.$$

Proof. We first show

$$(6.13) \quad p_i^\perp \cdot D \begin{pmatrix} \theta_1, \dots, \theta_d \\ a_1, \dots, a_d \end{pmatrix} = \sum_{j=1}^d D \begin{pmatrix} \theta_1, \dots, \theta_j, \dots, \theta_d \\ a_1, \dots, a_j + i, \dots, a_d \end{pmatrix}.$$

As $p_i^\perp h_j = h_{j-i}$ ([40], Chap. I, 5, Example 3), the action of $p_i^\perp$ on the column vector

$$\mathbf{c}((1-\zeta)^c \zeta^{-\theta} \varphi) = c_0 \cdot \mathbf{c} \left((1-\zeta)^c \zeta^{-\theta} \left(\sum_{i=0}^{n-1} h_i (1-\zeta)^i \right) \right)$$

is expressed as

$$p_i^\perp \cdot \mathbf{c}((1-\zeta)^c \zeta^{-\theta} \varphi) = \mathbf{c}((1-\zeta)^{a+i} \zeta^{-\theta} \varphi).$$

This relation and the ‘Leibniz rule’ $p_i^\perp(fg) = (p_i^\perp f)g + f(p_i^\perp g)$ imply the desired equation. (6.12) is obtained from (6.13) by using (4.22) as follows:

$$\begin{aligned} \kappa_d(p_i)^\perp \cdot D \begin{pmatrix} \theta_1, \dots, \theta_d \\ a_1, \dots, a_d \end{pmatrix} &= \sum_{j=1}^d \sum_{m=0}^i (-1)^m \binom{i}{m} D \begin{pmatrix} \theta_1, \dots, \theta_j, \dots, \theta_d \\ a_1, \dots, a_j + m, \dots, a_d \end{pmatrix} \\ &= \sum_{j=1}^d D \begin{pmatrix} \theta_1, \dots, \theta_j - i, \dots, \theta_d \\ a_1, \dots, a_j, \dots, a_d \end{pmatrix}. \end{aligned}$$

□

6.3.2. Boson-Fermion correspondence. To prove Proposition 6.8, we use the *Boson-Fermion correspondence*. Here we review some basic facts about it without proof. For details, see [18, 42].

Let $\mathcal{M} := \{M = (m_0, m_1, m_2, \dots) \mid m_0 > m_1 > \dots, m_j = -j \ (j \gg 1)\}$. Let v_m be infinitely many linearly independent vectors indexed by $m \in \mathbb{Z}$. Let

$$\mathcal{F} = \bigoplus_{M \in \mathcal{M}} \mathbb{C}v_M, \quad v_M := v_{m_0} \wedge v_{m_1} \wedge \dots$$

be the *Fermion-Fock space*. The vector $\Omega := v_0 \wedge v_{-1} \wedge v_{-2} \wedge \dots$ is called the *vacuum vector* of $\mathcal{F}$. For $m \in \mathbb{Z}$, $m \neq 0$, define $\alpha_m \in \text{End}_{\mathbb{C}} \mathcal{F}$ by the formula:

$$\alpha_m(v_{m_0} \wedge v_{m_1} \wedge \dots) = \sum_{j=0}^{\infty} v_{m_0} \wedge \dots \wedge v_{m_{j-1}} \wedge v_{m_j - m} \wedge v_{m_{j+1}} \wedge \dots$$

Then we have the *Heisenberg relation*:

$$[\alpha_m, \alpha_n] = m\delta_{m+n,0}.$$

There uniquely exists a linear isomorphism $\phi : \mathcal{F} \rightarrow \Lambda$ with the following properties:

$$\phi(\Omega) = 1, \quad \phi(\alpha_{-m}v) = p_m \phi(v), \quad \phi(\alpha_m v) = p_m^\perp \phi(v), \quad m \geq 1, \quad v \in \mathcal{F}.$$

Proposition 6.10. (see [42, §9.3], [18, §6]). *We have $\phi(v_{m_0} \wedge v_{m_1} \wedge \dots) = s_\lambda$, where $\lambda = (m_0, m_1 + 1, m_2 + 2, \dots)$ considered as a partition.*

6.3.3. *Proof of Proposition 6.8.*

Proof. Consider the subset $\mathcal{M}_d \subset \mathcal{M}$ which is defined by

$$\mathcal{M}_d = \{(m_j) \in \mathcal{M} \mid m_j = -j \ (j \geq d)\}$$

and the subspaces $\mathcal{F}_d, \mathcal{W}_d$ of $\mathcal{F}$ defined by

$$\mathcal{F}_d = \bigoplus_{M \in \mathcal{M}_d} \mathbb{C} \cdot v_M, \quad \mathcal{W}_d = \bigoplus_{M \in \mathcal{M} \setminus \mathcal{M}_d} \mathbb{C} \cdot v_M.$$

The space $\mathcal{F}$ decomposes as $\mathcal{F} = \mathcal{F}_d \oplus \mathcal{W}_d$. Let $v \mapsto \bar{v}, \mathcal{F} \rightarrow \mathcal{F}_d$ be the projection.

Let $\iota : \mathcal{F}_d \rightarrow \Lambda$ be the linear map

$$v_{m_0} \wedge \cdots \wedge v_{m_{d-1}} \wedge v_{-d} \wedge v_{-d-1} \wedge \cdots \mapsto D(-m_{d-1}, \dots, -m_0).$$

Note that $\iota(\Omega) = D(d-1, d-2, \dots, 0) = g_{R_d}$. Lemma 6.9 can be rewritten as

$$(6.14) \quad \kappa_d(p_i)^\perp \cdot \iota(v) = \iota\left(\overline{\alpha_{-i}(v)}\right), \quad v \in \mathcal{F}_d.$$

For $v \in \mathcal{F}_d$ define $\hat{s}_\lambda(v) = \overline{s_\lambda \cdot v} \in \mathcal{F}_d$, where the Schur function s_λ acts on $\mathcal{F}$ via the identification $p_m \mapsto \alpha_{-m}$ ($m \geq 1$). Accordingly we have

$$(6.15) \quad \kappa_d(s_\lambda)^\perp \cdot \iota(v) = \iota(\hat{s}_\lambda(v)), \quad v \in \mathcal{F}_d.$$

For a partition λ with $\ell(\lambda) \leq d$ we have the equation

$$\begin{aligned} & \hat{s}_\lambda \cdot \Omega \\ &= \phi^{-1}(s_\lambda \cdot 1) \\ &= v_{\lambda_1} \wedge v_{\lambda_2-1} \wedge \cdots \wedge v_{\lambda_d-d+1} \wedge v_{-d} \wedge v_{-d-1} \wedge \cdots \quad (\text{Prop. 6.10}) \\ &= v_{i_d-d} \wedge v_{i_{d-1}-d} \wedge \cdots \wedge v_{i_1-d} \wedge v_{-d} \wedge v_{-d-1} \wedge \cdots \quad (\text{by (6.6)}), \end{aligned}$$

and hence by substituting $v = \Omega$ to (6.15) we have

$$\kappa_d(s_\lambda)^\perp \cdot g_{R_d} = \kappa_d(s_\lambda)^\perp \cdot \iota(\Omega) = \iota(\hat{s}_\lambda \cdot \Omega) = D(i_1 - d, \dots, i_d - d).$$

□

7. DISCUSSION OF CONJECTURE 1.7

The aim of this section to explain some details about Conjecture 1.7 for the image of the quantum Grothendieck polynomials by Φ associated with arbitrary permutations.

7.1. λ -map. Recall that we set $k = n - 1$. Let $\mathcal{B}_k$ denote the set of k -bounded partitions. We recall the definition of a map $\lambda : S_n \rightarrow \mathcal{B}_k$ due to Lam and Shimozono [31, §6]. For $0 \leq i \leq n - 2$, let c_i denote the cyclic permutation $(i + 1, i + 2, \dots, n)$, and C denote the cyclic subgroup generated by $c_0 = (12 \cdots n)$. For $w \in S_n$, let $\tilde{w}$ be the unique element in the coset $C \cdot w$ such that $\tilde{w}(1) = 1$. There is a unique sequence $(m_1, \dots, m_{n-2})$ of non-negative integers such that

$$(7.16) \quad \tilde{w} = c_1^{m_1} c_2^{m_2} \cdots c_{n-2}^{m_{n-2}} \quad (0 \leq m_i \leq k - i).$$

Define $\lambda(w) = (1^{m_1} 2^{m_2} \cdots (n - 2)^{m_{n-2}})$, the partition whose multiplicity of i ($1 \leq i \leq n - 2$) is m_i .

Lemma 7.1. *Let μ be a partition contained in R_d . We have $\lambda(w_{\mu,d})^{\omega_k} = \mu^\vee$.*

Proof. We write $w = w_{\mu,d}$. We first assume that $w(1) = 1$, that is $\mu_d = 0$. It is straightforward to see $w = c_1^{m_1} c_2^{m_2} \dots c_{d-1}^{m_{d-1}} c_d^{m_d}$ with $m_i = w(i+1) - w(i) - 1$ ($1 \leq i \leq d-1$), and $m_d = n - w(d)$. Note that the Young diagram of $\lambda(w) = (1^{m_1} \dots d^{m_d})$ is contained in the conjugate of R_d , that is R_{n-d} . If we consider the complement $\mu^c = R_d \setminus \mu$ of μ in the rectangle R_d , the number of columns in μ^c having i boxes is m_i . The diagram of $\mu^\vee$ is obtained from μ^c by a rotation of 180 degrees. Now the conjugate of $\mu^\vee$ is nothing but $\lambda(w)$. Since the k -conjugate of a partition contained in R_d is the ordinary conjugate of it ([33, Remark 10]), the lemma follows in this case. If $w(i) > 1$, then $\tilde{w} = c_0^{-w(1)+1} w$ is $(d+l)$ -Grassmannian with some l such that $1 \leq l \leq k-d$. Let $\tilde{\mu} \subset R_{d+l}$ be the corresponding partition. One can check that $\tilde{\mu}^c = R_{d+l} \setminus \tilde{\mu}$ has the same shape as $\mu^c = R_d \setminus \mu$, and hence the proof is reduced to the case when $w(1) = 1$. $\square$

A k -bounded partition μ is k -irreducible if there is no k -rectangle R_d such that $\mu = R_d \cup \nu$ ($\nu \in \mathcal{B}_k$). This is equivalent to the inequalities $0 \leq m_i \leq k-i$ ($1 \leq i \leq n-2$), where m_i is the multiplicity of i in μ . Accordingly the image of the map λ is contained in the set $\mathcal{B}_k^*$ of all irreducible k -bounded partitions. In fact, one can easily see that $\mathcal{B}_k^*$ coincides the image of the λ -map.

Let S_n^* denote the set of permutations w in S_n such that $w(1) = 1$. The set S_n^* is a complete representatives of the coset space $C \setminus S_n$. In particular, the cardinality of $\mathcal{B}_k^*$ is $(n-1)!$

Recall that there is a remarkable involution $\omega_k : \mathcal{B}_k \rightarrow \mathcal{B}_k$, $\mu \mapsto \mu^{\omega_k}$ due to Lapointe and Morse [33]. One of the important properties is $\omega(s_\mu^{(k)}) = s_{\mu^{\omega_k}}^{(k)}$ [34, Theorem 38] where ω is the involution on Λ sending s_λ to $s_{\lambda'}$. One can see that ω_k preserves $\mathcal{B}_k^*$.

Example 7.2. The following tables give $\lambda(w)$ and its k -conjugate for $w \in S_n^*$ for $n = 4, 5$. The asterisk sign indicates that the permutation is not Grassmannian.

w	$\lambda(w)$	$\lambda(w)^{\omega_3}$
1234	$\emptyset$	$\emptyset$
1243	(2)	(1, 1)
1324	(2, 1)	(2, 1)
1342	(1)	(1)
1423	(1, 1)	(2)
1432*	(2, 1, 1)	(2, 1, 1)

w	$\lambda(w)$	$\lambda(w)^{\omega_4}$
12345	$\emptyset$	$\emptyset$
12354	(3)	(1, 1, 1)
12435	(3, 2)	(2, 2, 1)
12453	(2)	(1, 1)
12534	(2, 2)	(2, 2)
12543*	(3, 2, 2)	(2, 2, 1, 1, 1)
13245	(2, 2, 1)	(3, 2)
13254*	(3, 2, 2, 1)	(3, 2, 1, 1, 1)
13425	(3, 1)	(2, 1, 1)
13452	(1)	(1)
13524	(2, 1)	(2, 1)
13542*	(3, 2, 1)	(2, 2, 1, 1)
14235	(2, 1, 1)	(3, 1)
14253*	(3, 2, 1, 1)	(3, 2, 1, 1)
14325*	(3, 2, 2, 1, 1)	(3, 2, 2, 1, 1)
14352*	(2, 2, 1, 1)	(3, 2, 1)
14523	(1, 1)	(2)
14532*	(3, 1, 1)	(2, 1, 1, 1)
15234	(1, 1, 1)	(3)
15243*	(3, 1, 1, 1)	(3, 1, 1, 1)
15324*	(3, 2, 1, 1, 1)	(3, 2, 2, 1)
15342*	(2, 1, 1, 1)	(3, 1, 1)
15423*	(2, 2, 1, 1, 1)	(3, 2, 2)
15432*	(3, 2, 2, 1, 1, 1)	(3, 2, 2, 1, 1, 1)

7.2. Examples.

Example 7.3. Let $n = 5$. We first note some elements of $\tilde{g}_w$ are factored into a product of dual stable Grothendieck polynomials:

w	$\tilde{g}_w$
12543	$g_{1,1,1} \cdot g_{2,2}$
13254	$g_{1,1,1} \cdot g_{3,2}$
14532	$g_{1,1,1} \cdot g_2$
15243	$g_{1,1,1} \cdot g_3$
15324	$g_{2,2,1} \cdot g_3$
15342	$g_{1,1} \cdot g_3$
15423	$g_{2,2} \cdot g_3$
15432	$g_{1,1,1} \cdot g_{2,2} \cdot g_3$

Using Sage, we obtain the following expansion of $\tilde{g}_w$ in terms of K -theoretic k -Schur functions. Here we only write the non-Grassmannian elements in S_5^* .

$$\begin{aligned}
\tilde{g}_{12543} &= g_{2,2,1,1,1}^{(4)} - g_{2,2,1,1}^{(4)}, \\
\tilde{g}_{13254} &= g_{3,2,1,1,1}^{(4)} - g_{3,2,1,1}^{(4)}, \\
\tilde{g}_{13542} &= g_{2,2,1,1}^{(4)} - g_{2,2,1}^{(4)}, \\
\tilde{g}_{14253} &= g_{3,2,1,1}^{(4)}, \\
\tilde{g}_{14325} &= g_{3,2,2,1,1}^{(4)} - g_{3,2,1,1,1}^{(4)} - g_{3,2,2,1}^{(4)} + g_{3,2,1,1}^{(4)}, \\
\tilde{g}_{14352} &= g_{3,2,1}^{(4)} - g_{3,2}^{(4)}, \\
\tilde{g}_{14532} &= g_{2,1,1,1}^{(4)} - g_{2,1,1}^{(4)}, \\
\tilde{g}_{15243} &= g_{3,1,1,1}^{(4)}, \\
\tilde{g}_{15324} &= g_{3,2,2,1}^{(4)} - g_{3,2,1,1}^{(4)}, \\
\tilde{g}_{15342} &= g_{3,1,1}^{(4)} - g_{3,1}^{(4)}, \\
\tilde{g}_{15423} &= g_{3,2,2}^{(4)} - g_{3,2,1}^{(4)}, \\
\tilde{g}_{15432} &= g_{3,2,2,1,1,1}^{(4)} - g_{3,2,1,1,1,1}^{(4)} - 2g_{3,2,2,1,1}^{(4)} - g_{4,2,2,1}^{(4)} + g_{3,2,1,1,1,1}^{(4)} + g_{3,2,2,1}^{(4)} - g_{3,2,1,1}^{(4)}.
\end{aligned}$$

We finally provide one conjecture below.

Conjecture 7.4. Let $w_0 \in S_n$ denote the longest element. Then

$$\tilde{g}_{w_0} = \prod_{i=1}^{n-2} g_{(n-1-i)^i}.$$

Acknowledgements. We are informed that a general formulation of K -theoretic Peterson isomorphism valid for a simple simply connected G is proposed by Thomas Lam, Changzheng Li, Leonardo Mihalcea, and Mark Shimozono in [26]. We express thanks to them for communicating their work to us and for valuable discussions. We are grateful to Michael Finkelberg for valuable discussions. He kindly showed us his interpretation of τ -functions in terms of the Zastava space. During the preparation of the paper we also benefitted from the helpful discussions with and comments of many people, including Anders Buch, Rei Inoue, Hiroshi Iritani, Bumsig Kim, Cristian Lenart, Jennifer Morse, Satoshi Naito, Hiraku Nakajima, Hiroshi Naruse, Kyo Nishiyama, Masatoshi Noumi, Kaisa Taipale, Kanehisa Takasaki, Motoki Takigiku, Vijay Ravikumar, and Chris Woodward. We also thank Andrea Brini who drew

our attention to [24]. Special thanks are due to Tomoo Matsumura for showing us the proof of Proposition 5.1. In order to discover and check Conjecture 2, we used the open source mathematical software Sage [1]. The work was supported by JSPS KAKENHI [grant numbers 15K04832 to T.I., 26800062 to S.I., 16K05083 to T.M.].

REFERENCES

- [1] Sage Mathematics Software, version 7.5.1. <http://www.sagemath.org>, 2017.
- [2] Bezrukavnikov, R., M. Finkelberg, and I. Mirković. “Equivariant homology and K -theory of affine Grassmannians and Toda lattices.” *Compositio Mathematica* 141, no. 3 (2005): 746–68.
- [3] Braverman, A. and M. Finkelberg. “Finite difference quantum Toda lattice via equivariant K -theory.” *Transformation Groups* 10, no. 3–4 (2005): 363–86.
- [4] Braverman, A. and M. Finkelberg. “Semi-infinite Schubert varieties and quantum K -theory of flag manifolds.” *Journal of American Mathematical Society* 27, no. 4 (2014): 1147–68.
- [5] Buch, A. “A Littlewood-Richardson rule for the K -theory of Grassmannians.” *Acta Mathematica* 189, no. 1 (2002): 37–78.
- [6] Buch, A. “Combinatorial K -theory”, pp. 87–104. In *Topics in Combinatorial Studies of Algebraic Varieties.*, Trends in Mathematics, Birkhäuser Verlag Basel/Switzerland, 2005.
- [7] Buch, A., P.-E. Chaput, L. Mihalcea, and N. Perrin. “Finiteness of cominusule quantum K -theory.” *Annales scientifique de l’École normale supérieure* 46, no.3 (2013): 477–494.
- [8] Buch, A., P.-E. Chaput, L. Mihalcea, and N. Perrin. “Rational connectedness implies Finiteness of quantum K -theory.” *Asian Journal of Mathematics* 20, no.1 (2016): 117–22.
- [9] Buch, A. and L. Mihalcea. “Quantum K -theory of Grassmannians.” *Duke Mathematical Journal* 156 (2011): 501–38.
- [10] Fomin, S., S. Gelfand, and A. Postnikov. “Quantum Schubert polynomials.” *Journal of American Mathematical Society* 10 (1997): 565–96.
- [11] Gessel, I. M. and X. G. Viennot. “Binomial Determinants, Paths, and Hook Length Formulae.” *Advances in Mathematics* 58 (1985): 300–21.
- [12] Givental, A. “Homological geometry I: Projective hypersurfaces.” *Selecta Mathematica, New Series* 1 (1995): 325–45.
- [13] Givental, A. “Equivariant Gromov-Witten invariants.” *International Mathematics Research Notices* 1996, no. 13 (1996): 613–63.
- [14] Givental, A. “On the WDVV-equation in quantum K -theory.” *Michigan Mathematical Journal* 48 (2000): 295–304.
- [15] Givental, A. and B. Kim. “Quantum cohomology of flag manifolds and Toda lattices.” *Communications in Mathematical Physics* 168, no. 3 (1995): 609–41.
- [16] Givental, A. and Y.-P. Lee. “Quantum K -theory on flag manifolds, finite-difference Toda lattices and quantum groups.” *Inventiones mathematicae* 151, no. 1 (2003): 193–219.
- [17] Ikeda, T. and T. Shimazaki. “A proof of K -theoretic Littlewood-Richardson rules by Bender-Knuth-type involutions.” *Mathematical Research Letters* 21, no. 2 (2014): 333–39.
- [18] Iritani et al.(2015)] Iritani, H., T. Milanov, and V. Tonita. “Reconstruction and convergence in quantum K -theory via difference equations.” *International Mathematics Research Notices* 2015, no. 11 (2015): 2887–937.
- [19] Kac, V. G., A. K. Raina, and N. Rozhkovskaya. *Bombay Lectures on Highest Weight Representations of Infinite Dimensional Lie Algebras, 2nd edition*, Advanced Series in Mathematical Physics, vol.29. World Scientific, 2014.
- [20] Kashiwara, M. “The flag manifold of Kac-Moody Lie algebra.” In *Algebraic Analysis, geometry, and number theory (Baltimore, MD)*, pp. 161–190. Johns Hopkins Univ. Press, 1988.
- [21] Kim, B. “Quantum cohomology of flag manifolds G/B and quantum Toda lattices.” *Annals of Mathematics* 149 (1999): 129–48.
- [22] Kirillov, A. N. and T. Maeno. “A note on quantum K -theory of flag varieties and some quadric algebras.” *in preparation*.
- [23] Kostant, B. “The Solution to a Generalized Toda Lattice and Representation Theory.” *Advances in Mathematics* 34 (1979): 195–338.
- [24] Kostant, B. and S. Kumar. “ T -equivariant K -theory of generalized flag varieties.” *Journal of Differential Geometry* 32, no. 2 (1990): 549–603.
- [25] Kruglinskaya, O. and A. Marshakov. “On Lie Groups and Toda Lattice.” *Journal of Physics A* 48, no. 12 (2015): 125201, 26 pp.

- [25] Lam, T., L. Lapointe, J. Morse, A. Schilling, M. Shimozono, and M. Zabrocki. *k-Schur Functions and Affine Schubert Calculus*. Fields Institute Monographs. Springer, 2014.
- [26] Lam, T., C. Li, L. Mihalcea, and M. Shimozono. *in preparation*.
- [27] Lam, T. and P. Pylyavskyy. “Combinatorial Hopf algebras and K -homology of Grassmannians.” *International Mathematics Research Notices* 2007, no. 24 (2007) Art. ID rnm125, 48 pp.
- [28] Lam, T., A. Schilling, and M. Shimozono. “ K -theory Schubert calculus of the affine Grassmannian.” *Compositio Mathematica* 146, no. 4 (2010): 811–52.
- [29] Lam, T. and M. Shimozono. “From double quantum Schubert polynomials to k -double Schur functions via the Toda lattice.” <https://arxiv.org/abs/1109.2193>.
- [30] Lam, T. and M. Shimozono. “Quantum cohomology of G/P and homology of affine Grassmannian.” *Acta Mathematica* 204, no. 1 (2010): 49–90.
- [31] Lam, T. and M. Shimozono. “From quantum Schubert polynomials to k -Schur functions via the Toda lattice.” *Mathematical Research Letters* 19 (2012): 81–93.
- [32] Lapointe, L., A. Lascoux, and J. Morse. “Tableau atoms and a new Macdonald positivity conjecture.” *Duke Mathematical Journal* 116, no. 1 (2003): 103–46.
- [33] Lapointe, L. and J. Morse. “Tableau on $k + 1$ -cores, reduced words for affine permutations, and k -Schur functions.” *Journal of Combinatorial Theory, Series A* 112 (2005): 44–81.
- [34] Lapointe, L. and J. Morse. “A k -tableau characterization of k -Schur functions.” *Advances in Mathematics* 213, no. 1 (2007): 183–204.
- [35] Lascoux, A. and H. Naruse. “Finite sum Cauchy identity for dual Grothendieck polynomials.” *Proceedings of Japan Academy, Series A* 90, no. 7 (2014): 87–91.
- [36] Lascoux, A. and M.-P. Schützenberger. “Structure de Hopf de l’anneau de cohomologie et de l’anneau de Grothendieck d’une variété de drapeaux.” *Comptes Rendus de l’Académie des Sciences – Série I – Mathématiques* 295, no. 11 (1982): 629–33.
- [37] Lee, Y.-P. “Quantum K -theory I: foundation, Quantum K -theory II: computation and open problems.” *Duke Mathematical Journal* 121, no. 3 (2004): 389–424.
- [38] Lenart, C. and T. Maeno. Quantum Grothendieck polynomials. <https://arxiv.org/abs/math/0608232>.
- [39] Lenart, C. and A. Postnikov. “Affine Weyl groups in K -theory and representation theory.” *International Mathematics Research Notices* 2007, no. 12 (2007) Art. ID rnm038, 65pp.
- [40] Macdonald, I. G. *Symmetric functions and Hall polynomials*, 2nd edition. Oxford University Press, 1998.
- [41] Mihalea, L. “On equivariant quantum cohomology of homogeneous spaces: Chevalley formulae and algorithm.” *Duke Mathematical Journal* 140, no. 2 (2007): 321–50.
- [42] Miwa, T., M. Jimbo, and E. Date. *Solitons: Differential equations, symmetries and infinite dimensional algebras*. Cambridge Tracts in Mathematics vol.135. Cambridge University Press, 2000.
- [43] Morse, J. “Combinatorics of the K -theory of affine Grassmannians.” *Advances in Mathematics* 229 (2012): 2950–84.
- [44] Nakagawa, J., M. Noumi, M. Shirakawa, and Y. Yamada. “Tableau representation for Macdonald’s ninth variation of Schur functions.” pp. 180–95. Physics and Combinatorics 2000, Proceedings of the Nagoya 2000 International Workshop. World Scientific, 2000.
- [45] Ruijsenaars, S. “Relativistic Toda systems.” *Communications in Mathematical Physics* 133, no. 2 (1990): 217–47.
- [46] Shimozono, M. and M. Zabrocki. “Stable Grothendieck symmetric functions and Ω -calculus” *unpublished* (2003).

TAKESHI IKEDA, DEPARTMENT OF APPLIED MATHEMATICS, OKAYAMA UNIVERSITY OF SCIENCE, 1-1 RIDAI-CHO KITA-KU OKAYAMA-SHI 700-0005, JAPAN

email address: ike@xmath.ous.ac.jp

SHINSUKE IWAO, COLLEGE OF SCIENCE AND ENGINEERING, AOYAMAGAKUIN UNIVERSITY, 5-10-1 FUCHINOBE, CYUŌ-KU, SAGAMIHARA-SHI, KANAGAWA 252-5258, JAPAN

email address: iwao@gem.aoyama.ac.jp

TOSHIKI MAENO, DEPARTMENT OF MATHEMATICS, MEIJO UNIVERSITY, 1 CHOME-501 SHIOGAMAGUCHI, TENPAKU WARD, NAGOYA-SHI, AICHI 468-0073, JAPAN

email address: tmaeno@meijo-u.ac.jp